

## SNR 1987A: OPENING THE FUTURE BY REACHING THE PAST

SANGWOOK PARK,<sup>1</sup> SVETOZAR A. ZHEKOV,<sup>2,3</sup> DAVID N. BURROWS,<sup>1</sup> AND RICHARD MCCRAY<sup>2</sup>

Received 2005 September 21; accepted 2005 October 14; published 2005 November 8

### ABSTRACT

We report an upturn in the soft X-ray light curve of SNR 1987A in late 2003 (~6200 days after the explosion), as observed with the *Chandra X-Ray Observatory*. Since early 2004, the rapid increase of the 0.5–2 keV band X-ray light curve can no longer be described by the exponential density distribution model with which we successfully fitted the data between 1990 and 2003. Around day 6200, we also find that the fractional contribution to the observed soft X-ray flux from the decelerated shock begins to exceed that of the fast shock and that the X-ray brightening becomes “global” rather than “spotty.” We interpret these results as evidence that the blast wave has reached the main body of the dense circumstellar material all around the inner ring. This interpretation is supported by other recent observations, including a deceleration of the radial expansion of the X-ray remnant, a significant upturn in the mid-IR intensities, and the prevalence of the optical hot spots around the entire inner ring, all of which occur at around day 6000. In contrast to the soft X-ray light curve, the hard-band (3–10 keV) X-ray light curve increases at a much lower rate, which is rather similar to the radio light curve. The hard X-ray emission may thus originate from the reverse shock where the radio emission is likely produced. Alternatively, the low increase rate of the hard X-rays may simply be a result of the continuous softening of the overall X-ray spectrum.

*Subject headings:* supernovae: general — supernovae: individual (SN 1987A) — supernova remnants — X-rays: general — X-rays: stars

### 1. INTRODUCTION

The inner ring of SN 1987A is believed to be a relic structure produced by the massive progenitor’s equatorial stellar winds. The optical and X-ray brightness of SNR 1987A were expected to increase dramatically when the blast wave shock front hit this high-density structure (Borkowski et al. 1997a, 1997b). The soft X-ray light curve of SN 1987A has been monitored on a regular basis since 1991, first by the *ROSAT* satellite (Hasinger et al. 1996) and then by the *Chandra X-Ray Observatory* (Burrows et al. 2000; Park et al. 2002; Park et al. 2004, hereafter P04). Up to day ~3000 (since the SN), the soft X-ray emission from SNR 1987A was faint and was linearly increasing (Burrows et al. 2000). The first optical hot spot emerged in 1997 (Pun et al. 1997), and subsequent soft X-ray observations by the *Chandra X-Ray Observatory* (beginning in 1999 October) found an upturn in the soft X-ray light curve (Burrows et al. 2000), which deviated from the earlier linear increase. Since then, the soft X-ray light curve has brightened dramatically (Park et al. 2002; P04). The rapid brightening of the soft X-ray light curve over the first 4 years of *Chandra* observations has been successfully described with a simple model that assumes a constant velocity shock propagating into an ambient medium whose density increases exponentially with radius (P04; Park et al. 2005).

The emergence of optical hot spots in the inner ring has been interpreted in terms of the interaction of the shock front with irregularities (i.e., protrusions or clumps of dense gas, possibly produced by instabilities) on the inner surface of the inner ring (Michael et al. 2000). In this picture, the decelerated forward shock entering the dense protrusions is also expected to produce significant soft X-ray emission, and indeed soft X-ray bright

spots have emerged at these *same* locations. The fraction of the shock front interacting with high-density gas should increase gradually through the inclusion of more and more protrusions until the entire inner ring is finally engulfed. Recent optical images obtained with the *Hubble Space Telescope* have indeed revealed spectacular developments of the optically bright spots all around the entire inner ring by the end of 2003 (e.g., plate 1 in McCray 2005).

Given the evidence for bright X-ray emission from the hot spots, it is somewhat surprising that the simple model of a blast wave interacting with an exponential density distribution has provided such a good fit to the soft X-ray light curve up to this point. We report here a significant upturn of the soft X-ray light curve at day ~6200, which cannot be fitted with the simple exponential density model.

### 2. OBSERVATIONS AND DATA REDUCTION

We have now performed a total of 11 monitoring observations of SNR 1987A with the Advanced CCD Imaging Spectrometer (ACIS) on board *Chandra*. We reduced the data following the methods described in our previous papers (Burrows et al. 2000; Park et al. 2002). In order to derive the X-ray fluxes, we performed spectral fits using the two-component plane-parallel shock model of P04, which provides a useful approximation of the complex temperature/velocity structure of the blast wave. The detailed description of the observations, data reduction, and the image/spectral analyses of SNR 1987A, in which we obtained the X-ray fluxes and images presented in this Letter, can be found elsewhere (S. Park et al. 2005, in preparation). We note here that we use an updated version of the XSPEC shock model appropriate for the nonequilibrium ionization plasma.<sup>4</sup> We also regenerated the ancillary response functions using the ACIS *caldb* version 3.00 for all 11 observations in order to consistently correct for the quantum ef-

<sup>1</sup> Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802; park@astro.psu.edu, dburrows@astro.psu.edu.

<sup>2</sup> Joint Institute for Laboratory Astrophysics, University of Colorado, Box 440, Boulder, CO 80309; zhekov@super.colorado.edu, dick@jila.colorado.edu.

<sup>3</sup> On leave from the Space Research Institute, Sofia, Bulgaria.

<sup>4</sup> The unpublished version of the updated model has been provided by K. Borkowski.

efficiency degradation of the ACIS over the  $\sim 5$  yr period of the observations. The updated X-ray fluxes for all 11 ACIS-S3 observations are presented in Table 1.

### 3. SOFT X-RAY LIGHT CURVES

The latest X-ray light curves of SNR 1987A are shown in Figure 1. As of 2005 July, the observed 0.5–2 keV band X-ray flux is  $F_x \sim 19.4 \times 10^{-13}$  ergs  $\text{cm}^{-2}$   $\text{s}^{-1}$  ( $L_x \sim 1.42 \times 10^{36}$  ergs  $\text{s}^{-1}$  at  $d = 50$  kpc, after correcting for  $N_H = 2.35 \times 10^{21}$   $\text{cm}^{-2}$ ), which is an order of magnitude brighter than it was in 2000 January and 2 orders of magnitude brighter than in 1992. Figure 1 shows that the simple “exponential” model of P04 (*short-dashed curve*) adequately fits the observed soft X-ray flux only until day  $\sim 6200$ . The observed soft X-ray flux at later times increases significantly faster than the model prediction and cannot be fitted by this model. It is perhaps because this simple model does not accurately describe the soft X-ray emission from the blast wave shock interacting with the more complex density profile. The effect of this complex shock structure may have recently become significant, as indicated by the development of the optically bright spots all around the inner ring.

The forward shock is expected to encounter an extremely steep increase in the ambient density as it interacts with the dense inner ring, which will result in a substantial deceleration of the overall shock velocity and a significant increase in the soft X-ray emission. We thus construct a more sophisticated model of the light curve that considers the fast and decelerated shock components separately. This model is motivated by the picture proposed in Zhekov et al. (2005), which reconciles the slow velocities inferred from X-ray line profiles with the much faster velocities obtained from the expansion of the X-ray remnant: i.e., the long-term expansion is dominated by the rapid shock in the low-density H II region while the line widths are dominated by the slower forward shock in the dense protrusions. Following the basic arguments of P04, the observed X-ray flux can be expressed as  $F_x \propto n_0^2 V_0 T_0^{-0.6} + n_r^2 V_r T_r^{-0.6}$ , where  $n_0$ ,  $V_0$ , and  $T_0$  are the density, volume, and electron temperature for the less dense H II region, respectively, while  $n_r$ ,  $V_r$ , and  $T_r$  are those for the shocked portion of the dense protrusions. As the blast wave enters the dense clumps,  $T_r = T_0(n_0/n_r)$  can be assumed for an electron-ion temperature equilibration. The total emitting volume is given by  $V_t = V_r + V_0$ , and  $f \equiv V_r/V_t$  is the volume filling factor of the dense clumps. The H II region is expected to have roughly constant density inside the inner ring, and the blast wave radius is therefore a power law with time (Chevalier 1982). For the case of SN 1987A, with  $\rho \propto t^{-3} v^{-9}$ , the radius of the forward shock can be written as  $R = R_0 + R'(t/t_0 - 1)^{2/3}$ , where  $R_0$  is the radius (in units of arcseconds) at which the blast wave started to interact with the H II region,  $t_0$  is the time at  $R = R_0$ , and  $R'$  is a coefficient related to the radial expansion rate of the shock. Given that the number of optical spots has increased rapidly in recent years, we assume that the protrusion filling factor  $f$  increases exponentially with radius to a value of unity at the radius of the inner ring. For simplicity, we assume  $V_0$  and  $V_r \propto R^3$ . The observed soft X-ray flux can then be expressed as

$$F_x(\tau) = F_0 R(\tau)^3 (\tau - 1)^{0.4} \times \left\{ 1 + (\eta^{2.6} - 1) \exp \left[ -\frac{R_r - R(\tau)}{D} \right] \right\},$$

TABLE 1  
X-RAY FLUX OF SNR 1987A

Age <sup>a</sup> (days)	Observed Flux <sup>b</sup> ( <i>ROSAT</i> ) (0.5–2 keV)	Age <sup>a</sup> (days)	Observed Flux <sup>b</sup> ( <i>Chandra</i> ) (0.5–2 keV)	Age <sup>a</sup> (days)	Observed Flux <sup>b</sup> ( <i>Chandra</i> ) (3–10 keV)
1215	<0.23	4711	1.61 ± 0.66	4711	0.84 ± 0.57
1448	0.07 ± 0.09	5038	2.40 ± 0.22	5038	0.92 ± 0.21
1645	0.15 ± 0.04	5176	2.71 ± 0.54	5176	1.22 ± 0.41
1872	0.19 ± 0.04	5407	3.55 ± 0.43	5407	1.20 ± 0.44
2258	0.27 ± 0.05	5561	4.19 ± 0.46	5561	1.49 ± 0.64
2408	0.32 ± 0.07	5791	5.62 ± 0.45	5791	1.82 ± 0.46
2715	0.33 ± 0.11	5980	6.44 ± 0.52	5980	1.95 ± 0.62
3013	0.41 ± 0.06	6157	7.73 ± 0.62	6157	2.38 ± 0.57
		6359	11.48 ± 0.69	6359	2.40 ± 0.60
		6533	16.29 ± 0.65	6533	2.80 ± 0.73
		6716	19.41 ± 0.97	6716	3.26 ± 0.68

<sup>a</sup> Days since the SN explosion.

<sup>b</sup> X-ray fluxes are in units of  $10^{-13}$  ergs  $\text{cm}^{-2}$   $\text{s}^{-1}$ . The *ROSAT* fluxes are taken from Hasinger et al. (1996).

where  $\tau \equiv t/t_0$ ,  $D$  is the characteristic *scale height* of the exponential filling factor,  $\eta \equiv n_r/n_0$ , and  $R_r$  is the radius of the inner ring. Based on the previous observations (P04 and references therein), we adopt  $R_0 = 0''.6$ ,  $t_0 \sim 1200$  days,  $R_r \sim 0''.83$ . If we assume an average shock velocity of  $v \sim 3000$   $\text{km s}^{-1}$  for the last several years, these results imply  $R' \sim 0''.07$ .

We fit the soft X-ray light curve using this model in terms of  $F_0$ ,  $\eta$ , and  $D$ . The best-fit model, displayed in Figure 1 (*solid curve*), gives  $F_0 = (0.2 \pm 0.02) \times 10^{-13}$  ergs  $\text{cm}^{-2}$   $\text{s}^{-1}$ ,  $\eta = 7.3 \pm 0.4$ , and  $D = 0''.018 \pm 0''.001$ , with  $\chi^2/\nu = 13.1/15$ . The recent upturn in the soft X-ray flux appears to be the result of a significant increase in the fraction of the shock front interacting with the irregular inner boundary of the dense inner ring. As might be expected, the protrusion component now dominates the flux (providing  $\sim 96\%$  of the total flux at day 6700), while providing only a small contribution at early times (e.g.,  $\sim 9\%$  of the total flux at day 3000, shortly before the first optical bright spot appeared). We note that the density ratio  $\eta$  is considerably lower than previous estimates (P04 and references therein). We attribute the low density ratio to the simplified diagnostics of our model, which can only treat the detailed density structure around the inner ring as an “average” value.

### 4. SOFT X-RAY IMAGES

Given the recent upturn of the soft X-ray light curve and the prevalence of the optical hot spots around the entire inner ring, the X-ray images might also reveal evidence for the brightening of the entire SNR. We therefore examined the spatial distribution of the soft X-ray brightening at two epochs. In Figure 2, we present the 0.5–2 keV band intensity ratios of SNR 1987A between two pairs of observations. In each panel, the gray scales are cut above the average value in order to emphasize flux increases. It is evident that the strong brightening occurred mostly in the northern rim and the southeastern bright spot between 2000 and 2002 (Fig. 2a). By contrast, the intensity increase between 2002 and 2005 (after day  $\sim 6000$ ) is nearly ubiquitous (Fig. 2b), suggesting that the entire ring is now increasing in X-ray brightness.

### 5. DISCUSSION

Days  $\sim 6000$ – $6200$  seem to be an important milestone in the evolution of SNR 1987A. This is the point after which the observed soft X-ray light curve starts to deviate considerably from




FIG. 1.—Composite light curves of SNR 1987A. The *ROSAT* data are taken from Hasinger et al. (1996). The radio fluxes were obtained with the Australian Telescope Compact Array (ATCA; provided by L. Staveley-Smith) and are arbitrarily scaled for the purpose of display. The solid curve is the best-fit model for the 0.5–2 keV band light curve (*ROSAT+Chandra*) as determined in this work (§ 3). The short-dashed curve is the best-fit model from P04, which is extrapolated after day 6000. The long-dashed curve is the linear model fitted to the *ROSAT* data only and then extrapolated until day 7000. The two dotted vertical lines mark day 3700 and 6000, respectively. The inset is a log scale presentation showing the details of the *ROSAT*, radio, and hard X-ray light curves.

the model prediction that had successfully described the data obtained over the previous ~13 yr. It is remarkable that this soft X-ray upturn is accompanied by several other notable observational events: (1) The blast wave shock kinematics obtained with our recent *Chandra* gratings observations indicated significantly slower velocities ( $v \sim 300\text{--}1700 \text{ km s}^{-1}$ ) for the X-ray-emitting hot gas than shock velocities previously estimated by X-ray and radio images ( $v \sim 3000\text{--}4000 \text{ km s}^{-1}$ ) (Zhekov et al. 2005). (2) The fractional contribution to the total observed X-ray flux from the decelerated shock (based on the two-shock spectral modeling; S. Park et al. 2005, in preparation) has steadily increased over the past 5 yr and became dominant after day ~6200 (Fig. 3). (3) Until day ~5800, the radial expansion rate of the X-ray remnant was  $v \sim 4000 \text{ km s}^{-1}$  (P04). The latest *Chandra* data show that the X-ray radial expansion rate decreases substantially to  $v \sim 1600 \text{ km s}^{-1}$  after day ~6200 (J. Racusin et al. 2005, in preparation). (4) The entire inner ring became dominated by the optical hot spots by day ~6000 (e.g., plate 1 in McCray 2005). (5) Recent mid-IR observations of SNR 1987A, most likely indicating dust emission in the inner ring, show a

remarkable brightening since day ~6000 (Bouchet et al. 2005). The optical/IR observations, X-ray spectral results, X-ray expansion measurements, soft X-ray images, and soft X-ray light curves are all consistent with our picture that the shock is now interacting with dense gas all around the inner ring.

On the other hand, the hard X-ray flux is increasing much less rapidly than the soft X-ray flux, although it is still steeper than the extrapolated *ROSAT* light curve (Fig. 1, *long-dashed curve*). We note that the hard X-ray light curve is similar to the radio light curves (Fig. 1). The radio emission likely originates from synchrotron emission from the shocked ejecta behind the reverse shock (Manchester et al. 2005). The similarity between the light curves in the radio and the hard X-ray emission suggests that the hard X-rays might also be produced behind the reverse shock rather than the decelerated forward shock front. However, the radio map does not show a clearly better correlation with the hard X-ray image than with the soft X-ray image (Fig. 4). Alternatively, the low rate of brightening in the hard X-ray light curve may simply result from the overall softening of the X-ray spectrum as an increasing fraction of




FIG. 2.—The 0.5–2 keV band intensity ratio of SNR 1987A. (a) 2002 December (day 5791) to 2000 December (day 5038) ratio, overlaid with contours of the 0.5–2 keV image taken in 2002 December. (b) 2005 July (day 6716) to 2002 December (day 5791) ratio, overlaid with contours of the 0.5–2 keV image taken on 2005 July.




FIG. 3.—Fractional contributions of the slow and fast shock components to the total 0.5–2.0 keV band flux, based on the spectral fits of S. Park et al. (2005, in preparation).




FIG. 4.—Gray-scale X-ray images of SNR 1987A (2005 July) overlaid with the ATCA 9 GHz radio contours (2005 June). (a) Soft X-ray (0.4–0.5 keV) vs. radio. (b) Hard X-ray (3–8 keV) vs. radio. Both soft and hard X-ray images contain similar photon statistics of  $\sim 650$ –750 counts. The radio images have been provided by B. Gaensler and L. Staveley-Smith.

the forward shock front decelerates. If the hard X-rays are truly originating from the same locations where the radio emission is produced, we may expect the hard X-ray and radio light curves to increase at the same rate in the future. Otherwise, the hard X-ray flux increase rate may continue to decline. As SNR 1987A continues to brighten, we may achieve sufficient photon statistics even in the hard X-ray band to distinguish

between thermal and nonthermal nature, which will be critical to unveil the origin of the hard X-ray emission: e.g., broad (a few times  $10^3$  km s $^{-1}$ ) emission lines may be detected if the hard X-rays originate from a thermal plasma behind the reverse shock.

As the blast wave now sweeps through the inner ring, the soft X-ray brightening of SNR 1987A will be more spectacular than ever and the X-ray spectrum may significantly change in coming years. In this new stage, SNR 1987A will provide a unique, unprecedented opportunity for “real time” studies of the evolution of both the forward and reverse shocks. The soft X-ray emission from the main body of the inner ring would also be useful to study the past history of the progenitor star by monitoring the progress of the forward shock through this material. Continuous multiwavelength monitoring of SNR 1987A, including in X-rays, will be essential for such studies.

The authors thank K. Borkowski for providing the unpublished version of the shock model. We also thank B. Gaensler and L. Staveley-Smith for providing unpublished radio images and fluxes taken with the Australian Telescope Compact Array. This work was supported in part by the Smithsonian Astrophysical Observatory under *Chandra* grant GO5-6073X.

#### REFERENCES

- Borkowski, K. J., Blondin, J. M., & McCray, R. 1997a, *ApJ*, 476, L31  
 ———. 1997b, *ApJ*, 477, 281  
 Bouchet, P., et al. 2005, *ApJ*, submitted  
 Burrows, D. N., et al. 2000, *ApJ*, 543, L149  
 Chevalier, R. A. 1982, *ApJ*, 258, 790  
 Hasinger, G., Aschenbach, B., & Trümper, J. 1996, *A&A*, 312, L9  
 Manchester, R. N., Gaensler, B. M., Staveley-Smith, V. L., Kesteven, M. J., & Tzioumis, A. K. 2005, *ApJ*, 628, L131  
 McCray, R. 2005, in *IAU Colloq. 192, Cosmic Explosions*, ed. J. M. Marcaide & K. W. Weiler (Berlin: Springer), 77  
 Michael, E., et al. 2000, *ApJ*, 542, L53  
 Park, S., Burrows, D. N., Garmire, G. P., Nousek, J. A., McCray, R., Michael, E., & Zhekov, S. 2002, *ApJ*, 567, 314  
 Park, S., Zhekov, S. A., Burrows, D. N., Garmire, G. P., & McCray, R. 2004, *ApJ*, 610, 275 (P04)  
 ———. 2005, *Adv. Space Res.*, 35, 991  
 Pun, C. S. J., Sonneborn, G., Bowers, C., Gull, T., Heap, S., Kimble, R., Maran, S., & Woodgate, B. 1997, *IAU Circ.* 6665  
 Zhekov, S. A., McCray, R., Borkowski, K. J., Burrows, D. N., & Park, S. 2005, *ApJ*, 628, L127