

THE FAR-ULTRAVIOLET SPECTRA OF TW HYDRAE. II. MODELS OF H₂ FLUORESCENCE IN A DISK¹

GREGORY J. HERCZEG, BRIAN E. WOOD, AND JEFFREY L. LINSKY

JILA, University of Colorado and NIST, Boulder, CO 80309–0440; gregoryh@origins.colorado.edu, woodb@marmoset.colorado.edu, jlinsky@jila.colorado.edu

JEFF A. VALENTI

Space Telescope Science Institute, Baltimore, MD 21218; valenti@stsci.edu

AND

CHRISTOPHER M. JOHNS-KRULL

Department of Physics and Astronomy, Rice University, 6100 Main Street, MS-108, Houston, TX 77005; cmj@rice.edu

Received 2003 June 8; accepted 2004 February 4

ABSTRACT

We measure the temperature of warm gas at planet-forming radii in the disk around the classical T Tauri star (CTTS) TW Hya by modeling the H₂ fluorescence observed in *Hubble Space Telescope* Space Telescope Imaging Spectrograph and *Far Ultraviolet Spectroscopic Explorer* spectra. Strong Ly α emission irradiates a warm disk surface within 2 AU of the central star and pumps certain excited levels of H₂. We simulate a one-dimensional plane-parallel atmosphere to estimate fluxes for the 140 observed H₂ emission lines and to reconstruct the Ly α emission profile incident upon the warm H₂. The excitation of H₂ can be determined from relative line strengths by measuring self-absorption in lines with low-energy lower levels, or by reconstructing the Ly α profile incident upon the warm H₂ using the total flux from a single upper level and the opacity in the pumping transition. Based on those diagnostics, we estimate that the warm disk surface has a column density of $\log N(\text{H}_2) = 18.5^{+1.2}_{-0.8}$, a temperature $T = 2500^{+700}_{-500}$ K, and a filling factor of H₂, as seen by the source of Ly α emission, of 0.25 ± 0.08 (all 2 σ error bars). TW Hya produces approximately $10^{-3} L_{\odot}$ in the FUV, about 85% of which is in the Ly α emission line. From the H I absorption observed in the Ly α emission, we infer that dust extinction in our line of sight to TW Hya is negligible.

Subject headings: accretion, accretion disks — circumstellar matter — line: identification — stars: individual (TW Hydrae) — stars: pre-main-sequence — ultraviolet: stars

1. INTRODUCTION

Classical T Tauri Stars (CTTSs) are roughly solar-mass pre-main-sequence (PMS) stars that are accreting gas from their circumstellar disks. The extent and mass of dust in disks have been determined from IR spectral energy distributions (SEDs) and imaging (see review by Zuckerman 2001). Cold gas at large radii is traced by observations of molecules such as CO and HCN (e.g., Aikawa et al. 2002). Warmer neutral and ionized gases are used as diagnostics of accretion (e.g., Gomez de Castro & Lamzin 1999; Johns-Krull & Gafford 2002). However, observing the gas in the disk close to the star, which is important for the formation and evolution of planets, has been difficult. Gas at these radii can induce planet migration (Goldreich & Tremaine 1979), dampen the eccentricity of terrestrial planets (Agnor & Ward 2002), and is necessary for accretion onto giant planets (Lissauer 1993).

In search of gas at planet-forming radii, Calvet et al. (1991) investigated the formation of CO overtone lines at 2.3 μm in disks around CTTSs. These models predict that CTTSs as a group should show both CO emission and absorption, while

Herbig Ae/Be stars should show CO in emission. Using high spectral resolution observations, Carr et al. (1993) showed that the CO overtone emission from the probable Herbig Ae/Be star WL 16 arises in the inner disk, confirming the prediction of Calvet et al. (1991). Najita et al. (1996) find the same result for the Herbig Ae/Be star 1548C27. However, broader searches at high spectral resolution for CO overtone emission or absorption from CTTSs generally find no evidence for contributions to these lines from the disks around these stars (Casali & Eiroa 1996; Johns-Krull & Valenti 2001a). On the other hand, Najita, Carr, & Mathieu (2003) find that CO fundamental emission near 4.6–4.9 μm is common among CTTSs. They suggest that overtone emission has not been detected at the observed CO temperatures (1100–1300 K), mainly because the overtone bands have much lower transition rates. However, CO is not expected to be the dominant gas constituent in the disks around CTTSs, and it is the major component of the gas, H₂, that we seek to probe here.

Previous observations of H₂ have not provided a clear picture of the gas in the disk. Thi et al. (2001) detected cold H₂ emission in the pure rotational $S(0)$ and $S(1)$ lines from the *Infrared Space Observatory* (ISO) observations of a large sample of CTTSs, Herbig Ae/Be stars, and debris-disk stars. However, in ground-based observations of the $S(1)$ and $S(2)$ lines, using much smaller apertures than ISO, Richter et al. (2001) and Sheret et al. (2003) did not detect any H₂ emission from some of the same sources, indicating that the emission is

¹ Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. This work is also based on data obtained for the *FUSE* Science Team by the NASA-CNES-CSA *FUSE* mission, operated by Johns Hopkins University.

probably extended. On the other hand, the emission in the H_2 1–0 $S(1)$ transition detected by Bary, Weintraub, & Kastner (2003) in Phoenix spectra of four T Tauri stars is narrow and centered at the radial velocity of the stars, presumably formed in low-velocity circumstellar gas.

While H_2 lines in the IR are typically weak and are observed against a strong dust continuum, fluorescent H_2 lines dominate the far-ultraviolet (FUV) spectrum of CTTs at wavelengths longward of $Ly\alpha$. Ardila et al. (2002a) found that narrow H_2 lines can be blueshifted by up to 20 km s^{-1} in *Hubble Space Telescope* (HST) Goddard High Resolution Spectrograph spectra of CTTs, observed with a $2'' \times 2''$ aperture. High-resolution echelle and long-slit spectroscopy with HST Space Telescope Imaging Spectrograph (STIS) is providing the critical data needed to identify the source of the H_2 emission. Observations of T Tau reveal two components of H_2 emission. Saucedo et al. (2003) and Walter et al. (2003) detect off-source UV H_2 emission that is pumped near line center of $Ly\alpha$ and most likely produced where stellar outflows shock the surrounding ambient molecular material. The same set of fluorescent H_2 lines are detected in low-excitation HH objects such as HH43 and HH47 (Schwartz 1983; Curiel et al. 1995), suggestive of a similar excitation mechanism. Walter et al. (2003) find that the on-source H_2 emission is photoexcited by a much broader $Ly\alpha$ emission line, likely produced by the accretion shock at the surface of T Tau.

In Herczeg et al. (2002, hereafter Paper I), we presented observations of H_2 fluorescence in the UV spectrum of TW Hya obtained with HST STIS and the *Far Ultraviolet Spectroscopic Explorer* (FUSE). Several lines of evidence suggested a disk origin for the H_2 emission: (i) the H_2 emission is not spatially extended beyond a point source in the cross-dispersion direction, (ii) the H_2 lines are emitted interior to TW Hya's wind because emission in one H_2 line is suppressed by C II wind absorption, (iii) the H_2 line centroids have the same radial velocity as the photospheric lines of TW Hya, (iv) no H_2 absorption is detected against the $Ly\alpha$ emission, and (v) the TW Hya Association is isolated from large reservoirs of interstellar molecular material, making a circumstellar origin for the H_2 emission unlikely. Because the H_2 emission is not extended beyond a point source, the spatial resolution of HST STIS restricts the emitting region to be within 2 AU of the star.

Analysis of on-source IR H_2 emission lines to date has been limited because typically only one or two lines are observed. The IR lines could be excited by shocks, UV fluorescence, X-rays, or thermal heating. In contrast, over 140 H_2 lines from 19 different upper levels are detected in the FUV spectrum of TW Hya. The excitation of H_2 can be determined from relative line strengths in this rich spectrum either by measuring self-absorption in low-excitation members of individual fluorescent progressions or by using an assumed $Ly\alpha$ profile to reconstruct initial level populations in various pumping transitions.

In this paper, we calculate the temperature of warm gas at planet-forming distances by modeling warm H_2 in a disk around TW Hya. We use the H_2 emission to reconstruct the $Ly\alpha$ profile incident on the fluoresced H_2 . The observed $Ly\alpha$ profile differs from the incident profile because of interstellar absorption along our line of sight to TW Hya. We discuss the strength of $Ly\alpha$ emission and its effect on the circumstellar disk.

2. PROPERTIES OF TW HYDRAE

TW Hya is the namesake of the 10 Myr old TW Hya Association (TWA), which most likely originated in the

Sco-Cen OB association (Mamajek, Lawson, & Feigelson 2000). TW Hya is one of the oldest known stars still in the CTT phase, and its mass accretion rate of about $(2-3) \times 10^{-9} M_\odot \text{ yr}^{-1}$ is lower than the mass accretion rate of most of the 1–2 Myr old CTTs that are found in Taurus (Kenyon & Hartmann 1995). From PMS evolutionary tracks, Webb et al. (1999) estimate that TW Hya has a mass of $0.7 M_\odot$ and a radius of $1.0 R_\odot$. Gaseous material is often associated with young stars but is not found in the nearby TWA, so the extinction is low. The TWA is located at a distance of approximately 56 pc (Webb et al. 1999), compared with 140 pc for Taurus. Consequently, TW Hya is by far the brightest known CTT in the UV and in X-rays. Most of the UV radiation from TW Hya is presumably produced at the accretion shock. The disk is viewed face-on (e.g., Zuckerman, Forveille, & Kastner 1995; Alencar & Batalha 2002), which prevents significant Keplerian broadening in the H_2 line profiles. The disk mass is roughly $(1.5-3) \times 10^{-2} M_\odot$ (Wilner et al. 2000; Trilling et al. 2001) and extends more than 225 AU from the star (e.g., Krist et al. 2000). Based on estimates of the stellar mass, radius, and rotation period of TW Hya, Johns-Krull & Valenti (2001a) calculate that the disk at $6 R_*$ corotates with the stellar surface. In the Shu et al. (1994) model, the disk is truncated at corotation. Weinberger et al. (2002) estimate from modeling the IR SED of TW Hya that the dust in the disk is truncated at about $11 R_*$. In their models of the IR SED, Calvet et al. (2002) found an underabundance of micron-sized dust grains within 4 AU of TW Hya. Figure 1 shows a schematic model of TW Hya and its disk.

Since the mass accretion rate of CTTs can vary, we calculate the mass accretion rate at the epoch when we observed TW Hya. We then proceed to estimate the extinction using H I absorption against $Ly\alpha$ emission and the SED of the NUV continuum that is produced in an accretion shock. The observations discussed in this paper were described in Paper I and are listed in Table 1.

2.1. Mass Accretion Rate

The strong excess blue and UV emission from CTTs can be modeled as emission from shocks at accretion footpoints




FIG. 1.—Schematic diagram of our model of TW Hya and its disk (not to scale). FUV radiation, primarily $Ly\alpha$, is produced at the accretion shock and heats the disk. The warm H_2 is located in surface layers of the disk within 2 AU of the star. The hot surface of the disk, composed of atoms and ions, probably extends inward to about $6 R_*$ (Johns-Krull & Valenti 2001a), where it is truncated by the stellar magnetosphere.

TABLE 1
LIST OF OBSERVATIONS

Star	Date	Instrument	Exposure Time (s)	Grating	Aperture (arcsec)	Wavelength Range (Å)
TW Hya	2000 May 7	<i>HST</i> STIS	260	G430L	$52. \times 0.2$	2900–5700
	2000 May 7	<i>HST</i> STIS	1675	E230M	0.2×0.2	2150–2950
	2000 May 7	<i>HST</i> STIS	2300	E140M	0.5×0.5	1160–1700
	2000 Jun 22	<i>FUSE</i>	2081	^a	$30. \times 30.$	900–1185
V819 Tau	2000 Aug 30	<i>HST</i> STIS	360	G430L	$52. \times 0.1$	2900–5700
	2000 Aug 31	<i>HST</i> STIS	2325	E140M	0.2×0.2	2300–3100
	2000 Aug 30	<i>HST</i> STIS	5700	E140M	0.2×0.06	1160–1700

^a The *FUSE* instrument consists of several channels.

on the star, which leads to estimates of the mass accretion rate. We estimate the accretion rate of TW Hya by following the method of Valenti, Basri, & Johns (1993), who modeled observed blue spectra with the sum of a photospheric template and emission from a slab of hot isothermal hydrogen to crudely simulate emission from the accretion shock.

We use the *HST* STIS spectrum of the K7 weak-lined TTS V819 Tau (see Table 1), spanning 1100–6000 Å, as a non-accreting template for the photospheric emission from TW Hya. For V819 Tau we adopt a visual extinction of $A_V = 1.52$, which is the average of values determined by Kenyon & Hartmann (1995) and White & Ghez (2001). We then use the interstellar extinction law of Cardelli, Clayton, & Mathis (1989) to remove the effects of extinction from the observed spectrum of V819 Tau. The NUV and FUV emission from V819 Tau is much fainter than that of TW Hya and is not significant in this analysis.

The spectrum of the model slab is determined by the slab temperature T , density n , thickness l , extinction A_V toward the slab, and a slab surface area parameter δ , as described in Valenti et al. (1993). The slab surface area parameter δ corresponds to the fraction of the stellar surface area covered by accretion-related emission and is calculated from the scale factor applied to the slab spectrum. The scaling factor between the nonaccreting template star and TW Hya is another free parameter. We find the best-fit model parameters by fitting the synthetic slab spectrum plus the scaled template spectrum to the NUV and optical spectrum of TW Hya, minimizing χ^2 using the *amoeba* function in IDL, which is based on the downhill simplex minimization method. The best-fit parameters for the slab are $A_V = 0.0$, $T = 9900$ K, $n = 1.7 \times 10^{14}$ cm⁻³, $l = 125$ km, $\delta = 0.004$, and a template scaling factor of 1.18. The total slab flux integrated over all wavelengths is 3.5×10^{-10} ergs cm⁻² s⁻¹.

Figure 2 shows our fit to the optical spectrum of TW Hya, after subtracting the scaled template spectrum of V819 Tau. The accretion continuum calculated in this model is consistent with the observed NUV continuum. However, the observed continuum flux rises shortward of 1700 Å, which could result from (i) an H₂ dissociation continuum produced by electron collisions (Liu & Dalgarno 1996), (ii) H₂ fluorescence pumped by the FUV continuum, or (iii) by a hot accretion or activity component. The SED of the FUV continuum is similar to the continuum observed in HH2, which is probably produced by H₂ dissociation (Raymond, Blair, & Long 1997). However, the FUV continuum of TW Hya appears smooth and does not show strong H₂ emission lines at 1054 and 1101 Å, which are detected toward HH2.

Assuming that half of the potential energy of the accreting gas is converted into hydrogen continuum emission,² then the mass accretion rate of TW Hya can be estimated by

$$\dot{M}_* = \frac{1.25 R_* L_{\text{slab}}}{GM_*}, \quad (1)$$

where $L_{\text{slab}} = 0.034 L_\odot$ is the luminosity of the slab. The leading factor of 1.25, derived by Gullbring et al. (1998) for a magnetospheric accretion geometry, replaces the factor of 2 from a spherical accretion geometry. We adopt 0.7 and 1.0 R_\odot for the mass and radius of TW Hya (Webb et al. 1999). Uncertainties in the stellar mass, radius, and distance are all of order 20%. We calculate an accretion rate onto the star of $\dot{M} \approx 2 \times 10^{-9} M_\odot \text{ yr}^{-1}$, which is consistent with the accretion rate of $2 \times 10^{-9} M_\odot \text{ yr}^{-1}$ estimated by Alencar & Batalha (2002) using a similar method and is larger than the

² See Lynden-Bell & Pringle (1974) and Hartigan et al. (1991) for details.




FIG. 2.—Solid line shows the observed spectrum of TW Hya, after subtracting the template spectrum of V819 Tau. Dashed line shows the synthetic spectrum of a pure hydrogen slab fit to the observed Balmer jump and the Balmer lines. The pure hydrogen continuum accurately approximates the NUV continuum of TW Hya but significantly underestimates the FUV continuum. The rise of the FUV continuum shortward of 1700 Å is indicative of an H₂ dissociation continuum produced by collisions with energetic electrons, although it could also be produced by H₂ fluorescence caused by FUV pumping or an additional accretion or activity component.




FIG. 3.—Ly α emission from TW Hya is partially absorbed by H I in the line of sight to TW Hya. The interstellar and circumstellar media, centered at about 0 km s⁻¹, account for the red side of the absorption. The blue side of the absorption is produced by H I in the wind. The dashed line shows our best fit to the observed Ly α profile (solid line) using two Gaussian emission profiles to model the intrinsic Ly α line before the absorption, while the dotted and dashed-dotted lines assume different H I column densities that poorly fit the red side of the observed profile.

accretion rate of $5 \times 10^{-10} M_{\odot} \text{ yr}^{-1}$ estimated by Muzerolle et al. (2000) using an analysis of the H α line.

2.2. Extinction to TW Hya

In this section, we argue that the extinction to TW Hya is negligible by measuring the hydrogen column density directly from the shape of the observed Ly α line and then using an ISM extinction law to infer the dust extinction. This argument does not necessarily imply that the extinction toward the warm H₂ is negligible, because some dust could be mixed with the warm H₂, leading to extinction of H₂ fluxes without any extinction of the stellar or accretion emission. We will discuss this possibility in § 5.

Figure 3 shows the observed Ly α profile, which is likely produced in an accretion column, contains a dark, broad absorption feature that extends from +200 to -500 km s⁻¹. The absorption is produced by H I in the wind and in interstellar and circumstellar material. Because circumstellar and interstellar absorption are indistinguishable in our data, we model these two features with a single absorption feature. All UV spectra of CTTSs show similar wind absorption in Mg II (Ardila et al. 2002b), so the star must be occulted by the outflow, regardless of inclination angle.

In the following analysis, we fit two Gaussian emission profiles and either one or two Voigt absorption profiles to the observed Ly α profile. The intrinsic emission profile is assumed to be a double-Gaussian profile, and the parameters of the Gaussians are allowed to vary to best fit the observed profile when combined with the absorption. Based on the widely separated wind and interstellar absorption features observed in the O I 1302 Å, C II 1334 Å, and Mg II 2795 Å lines, the interstellar and wind absorption components toward TW Hya must be centered at $v = 0$ and -130 km s⁻¹, respectively. For the interstellar component we assume a Doppler parameter of $b = 11 \text{ km s}^{-1}$, which corresponds to a temperature of 7000 K that is typical of the local (warm) ISM (Redfield & Linsky 2000). An interstellar column density is then determined by fitting the absorption longward of line

center, where any wind absorption should be negligible. The wind absorption extends to a velocity of at least -500 km s⁻¹ and likely has a large Doppler parameter owing to velocity gradients along our line of sight. Given these assumptions, the interstellar neutral hydrogen column density in our line of sight is $\log N(\text{H I}) = 19.05 \pm 0.15$ (Fig. 3). The upper limit of the column density in the wind is $N(\text{H I}) < 19.4$ for Doppler parameters $b > 25 \text{ km s}^{-1}$.

In a more conservative test to find the upper limit to the hydrogen column density, we fit the Ly α absorption with a single component, effectively combining any interstellar and wind absorption. We vary the central wavelength of the absorption and set a low Doppler parameter, so that the damping wings generate the width of the absorption profile. Figure 4 shows that the maximum possible column density consistent with the data is $\log N(\text{H I}) = 19.75$.

The conservative upper limit of the hydrogen column density toward TW Hya is therefore $\log N(\text{H}) < 19.75$. In the only previous measurement of $N(\text{H I})$ to TW Hya, Kastner et al. (1999) used X-ray spectral energy distributions from *ROSAT* and *ASCA* to estimate higher column densities of $\log N(\text{H I}) = 20.5 \pm 0.4$ and 21.4 ± 0.1 , respectively. However, the X-ray analyses are highly reliant on the accuracy of plasma codes. Kastner et al. (1999) estimate the hydrogen column density by assuming solar abundances and using a two-temperature emission measure rather than a differential emission measure distribution. However, in their analysis of the *Chandra* X-ray spectrum of TW Hya, Kastner et al. (2002) found that the abundances show an inverse-FIP effect. We further note that the sensitivity and flux calibration of *ROSAT* and *ASCA* are relatively poor at soft energies, where absorption from H I ionization is measured. Figure 4 shows that the absorption calculated for the Ly α line for $\log N(\text{H I}) = 20.5$ is significantly broader than the observed absorption profile. Our comparison of the calculated transmission function with the observed Ly α profile provides a much simpler and more direct test of hydrogen column density than the X-ray analyses. The discrepancy between the UV and X-ray measurements of $N(\text{H})$ could in principle be due to a variable circumstellar absorber, which Kastner et al. (1999) invoked to explain the order-of-magnitude discrepancy in calculations of $N(\text{H})$ from *ASCA* and *ROSAT*. However, the many ultraviolet observations of TW Hya by *IUE*, *HST* STIS, and *FUSE*, which are widely




FIG. 4.—Transmission percentage across the observed Ly α profile (solid line) for column densities $\log N(\text{H}) = 19.75$ (dashed line) and $\log N(\text{H}) = 20.5$ (dotted line), assuming no Doppler broadening. This assumption places a strong upper limit of $N(\text{H I}) < 19.75$ toward TW Hya.

spaced in time, show no evidence for large fluctuations in extinction.

We estimate an H_2 column density in our line of sight to TW Hya with *FUSE*, as the bandpass contains many H_2 transitions from low rotational levels of the ground electronic state, which are often seen in absorption owing to the ISM at temperatures typically ~ 100 K (Rachford et al. 2002). H_2 absorption lines from $v''=0$, $v''=0, 1, 2, 3, 5$ can in principle be observed against $O\text{ VI}$ emission (e.g., Roberge et al. 2001). The $O\text{ VI}$ profiles are noisy (see Fig. 5), but we estimate an upper limit of $\log N(H_2) < 18$ by calculating the transmission percentage for a range of temperatures and column densities, meaning the molecular fraction of H toward TW Hya is very low.

When we convert total hydrogen column density, $N(H\text{ I}) + 2N(H_2) = N(H)$, to a reddening using the interstellar relationship of Bohlin, Savage, & Drake (1978),

$$\frac{N(H\text{ I} + H_2)}{E(B-V)} = 5.8 \times 10^{21} \text{ cm}^{-2}, \quad (2)$$

we find that the reddening toward TW Hya is $E(B-V) = A_V/R_V < 0.01$, where R_V is a dimensionless parameter that ranges between 2.5 and 6, depending on the size distribution of dust grains. Significant deviations from the assumed dust-to-gas ratio can occur in star-forming regions such as ρ Oph, in which the dust grains are likely larger than they are in the ISM. However, the low molecular fraction toward TW Hya is consistent with a low extinction (Savage et al. 1977; Rachford et al. 2002) and is not consistent with material found toward star-forming regions. We conclude that this relationship provides a reasonable estimate of the extinction along our sight line to TW Hya. Even if the dust-to-gas ratio along the line of sight to TW Hya were for some reason enhanced by an order




FIG. 5.— $O\text{ VI}$ emission line from TW Hya (solid line), as observed with the LiF1a detector on *FUSE*. We place a limit of $N(H_2) < 18$ in our line of sight toward TW Hya by comparing the observed emission with the simulated transmission percentage given the resolution of *FUSE*, through a medium of H_2 with $T = 100$ K and $N(H_2) = 17.5$ (dashed line) and $N(H_2) = 19.0$ (dotted line). We would observe many H_2 absorption lines against the $O\text{ VI}$ emission from TW Hya for hotter excitation temperatures of H_2 .




FIG. 6.—NUV continuum (solid line) is well fitted by the accretion model (dashed line) for $A_V = 0.0$, but if $A_V = 0.5$, the rise in the continuum below 2600 Å is not fitted well.

of magnitude, then dust extinction would reduce the brightness of TW Hya by a factor of 2 and have only a 20% effect on the relative fluxes between 1200 and 1650 Å.

Inspection of the NUV continuum (Fig. 6) confirms the small extinction to TW Hya. If we assume that the NUV continuum is produced in an isothermal slab, as modeled in § 2.1, then it can be well fitted with a hydrogen continuum only for extinctions $A_V < 0.2$. As noted in § 2.1, an accretion continuum best fits the NUV continuum for $A_V = 0.0$.

3. MODELING H_2 LINE FLUXES

In Paper I, we measured a total line emission flux of 1.94×10^{-12} ergs $s^{-1} \text{ cm}^{-2}$ by summing 146 H_2 emission lines in the 19 progressions (the set of R and P transitions from a common upper level) listed in Table 2. Most of these lines are Lyman-band transitions fluoresced by the strong, broad $Ly\alpha$ emission line.

Wood, Karovska, & Raymond (2002) discovered in their analysis of *HST* STIS observations of Mira B that H_2 fluxes weaken toward short wavelengths because of increasing H_2 line opacities. This occurs because shorter wavelength lines are transitions to low energy levels, which are more populated than higher levels. Figure 7 demonstrates that the same effect occurs in the H_2 fluorescence sequences detected toward TW Hya. In most sequences, fluxes in many H_2 emission lines at short wavelengths are weaker than predicted from theoretical branching ratios because of significant absorption in the same transitions. Transitions from levels with low excitation energies, such as $v' = 0$, $J' = 1$, have significant optical depth below 1300 Å, but this does not happen for transitions from levels with large excitation energies such as $v' = 4$, $J' = 18$ (see Fig. 7). The $v' = 4$, $J' = 18$ level does not have any downward transitions with significant opacity in the STIS bandpass.

Following the Monte Carlo method of Wood et al. (2002), we compute H_2 opacities by modeling H_2 fluorescence in a one-dimensional isothermal plane-parallel atmosphere of H_2 irradiated by $Ly\alpha$ photons. We assume complete frequency and angular redistribution when photons scatter in the slab. Forty-one wavelength grid points across each H_2 line profile track differences in escape probability across the line profile because a photon that scatters into the wings of a line escapes from a slab more easily than a photon at line center. A $Ly\alpha$

TABLE 2
H₂ PUMPING TRANSITIONS

Pump	λ_{calc} (Å)	E''^a (eV)	A_{ul}^b (10^7 s^{-1})	d_{abs}^c (%)	$\text{EQW}_{\text{slab}}^d$ (Å)	R_{unseen}^e	P_{Dis}^f	$P_{\text{Dis, mod}}^g$	E_{dis}^h (eV)	M_{dis}^i	F_{mod}^j
1–1 $P(11)$	1212.425	1.36	6.6	0.23	0.067	0.22	1.4(–5)	0.0	0.07	0.0	4.5
3–1 $P(14)$	1213.356	1.79	10.1	0.43	0.028	0.38	0.015	0.018	0.10	2.5	5.9
4–2 $R(12)$	1213.677	1.93	3.9	0.50	0.0092	0.30	0.050	0.067	0.15	4.4	2.7
3–1 $R(15)$	1214.465	1.95	10.0	0.37	0.044	0.16	0.031	0.039	0.02	15.6	17
4–3 $P(5)$	1214.781	1.65	5.5	0.43	0.030	0.25	1.9(–3)	2.0(–3)	0.11	0.0	11
4–3 $R(6)$	1214.995	1.72	3.0	0.50	0.0092	0.68	9.2(–3)	0.011	0.07	0.5	2.4
1–2 $R(6)$	1215.726	1.28	13.6	0.26	0.062	0.06	2.4(–6)	0.0	0.08	0.0	16
1–2 $P(5)$	1216.070	1.20	15.9	0.17	0.075	0.07	4.8(–7)	0.0	0.06	0.0	33
3–3 $R(2)$	1217.031	1.50	0.40	0.50	0.0015	0.45	2.8(–4)	0.0	0.05	0.0	2.4
3–3 $P(1)$	1217.038	1.48	1.7	0.50	0.0030	0.02	1.3(–4)	0.0	0.06	0.0	2.9
0–2 $R(0)$	1217.205	1.01	6.6	0.39	0.0413	0.09	4.3(–9)	0.0	0.07	0.0	38
4–0 $P(19)$	1217.410	2.21	4.4	0.47	0.014	0.47	0.42	0.51	0.18	64.0	5.1
0–2 $R(1)$	1217.643	1.02	7.8	0.21	0.0685	0.23	5.2(–9)	0.0	0.07	0.0	32
2–1 $P(13)$	1217.904	1.64	9.3	0.28	0.059	0.09	1.7(–3)	2.9(–3)	0.02	1.3	18
3–0 $P(18)$	1217.982	2.03	3.2	0.50	0.0068	0.57	0.19	0.21	0.11	9.5	1.9
2–1 $R(14)$	1218.521	1.79	7.6	0.43	0.028	0.42	5.9(–3)	6.5(–3)	0.09	0.6	3.6
0–2 $R(2)$	1219.089	1.05	8.2	0.29	0.0576	0.47	6.6(–9)	0.0	0.06	0.0	2.1
0–2 $P(1)$	1219.368	1.02	20.1	0.28	0.0599	0.30	3.9(–9)	0.0	0.06	0.0	1.2
0–5 $P(18)$	1548.146	3.79	19.1	0.50	0.0	0.36	1.1(–4)	0.0	0.11	0.0	3.0

^a Lower energy level.

^b Radiative decay coefficient (Abgrall et al. 1993).

^c Absorption depth of Ly α photon.

^d Equivalent width of absorption profile within slab.

^e Correction for unseen lines from upper level.

^f Dissociation probability from upper state per electronic excitation, $a(b) = a \times 10^b$.

^g Dissociation probability from best-fit model.

^h Average kinetic energy released owing to H₂ dissociation.

ⁱ H₂ dissociation rate for best-fit model ($10^{-12} M_{\odot} \text{ yr}^{-1}$).

^j Ly α flux absorbed by lower level ($10^{-14} \text{ ergs cm}^{-2} \text{ s}^{-1}$).




FIG. 7.—For four different upper levels, we plot the normalized ratio of the observed de-excitations in each H₂ line divided by the respective branching ratio (circles). The squares show the fluxes predicted based on the optical depth in each lower level for our adopted model for $\log N(\text{H}_2) = 18.5$ and $T = 2500 \text{ K}$ (see § 5). The dotted and dashed lines show extinction curves for $A_V = 0.5$ and 1.5 , respectively. Emission in an optically thin medium, with no intervening extinction, would result in ratios of 1.0. The apparent wavelength dependence results from the larger optical depth in those levels of H₂, which have lower excitation energies. The H₂ line fluxes from $v' = 4$, $J' = 18$ show no wavelength dependence because the observed lines all have lower levels with energies large enough that they are not populated.

photon enters the slab at an angle θ with respect to the slab surface normal and photoexcites an H₂ molecule. The H₂ molecule quickly de-excites to the ground electronic state through one of the many available routes, or dissociates with a probability P_{Dis} (listed in Table 2, calculated by Abgrall, Roueff, & Drira 2000). Depending on the optical depth of H₂ in the lower level of the downward transition, the resulting photon may be temporarily absorbed by another H₂ molecule or may emerge from either side of the slab at an angle ϕ with respect to the normal. In our simple model, H₂ line photons created fluorescently in the slab either escape, are converted to another H₂ line in the same progression, or dissociate an H₂ molecule. No other photon processes are considered. We tabulate photons that exit the slab on the Ly α entry side (reflected photons) and opposite side (transmitted photons) using 40 bins of equal solid angle on each side of the slab. We refer the reader to Wood et al. (2002) for further details of this Monte Carlo code.

We use the term “model” to refer to the simulated plane-parallel atmosphere described in the preceding paragraph, and the term “geometry” to refer to the morphology of the H₂ around the Ly α emission source. In the context of the model, the H₂ line fluxes depend on the temperature and column density of the slab, the angle of incidence θ of the Ly α photon into the disk, the outward angle ϕ of the emerging photon, the side of the slab from which the photon escapes, and the optical depth of the transition. We assume that extinction is negligible in the slab and in any interstellar material. Figure 8 shows two possible geometries that we use to approximate a thick disk and a thin disk. The term “thick disk” applies to a star-disk geometry in which most of the H₂ illuminated by Ly α is at the




FIG. 8.—Two plausible geometries for H_2 fluorescence in the surface layers of a protoplanetary disk. The $Ly\alpha$ enters the surface layer of H_2 at angle θ with respect to the normal and is reprocessed into H_2 emission, which escapes the disk at angle ϕ .

inner edge of the disk. The term “thin disk” applies to a geometry in which most of the H_2 illuminated by $Ly\alpha$ is at the disk surface. In this section and § 4, we present detailed results for the thick disk geometry that reproduces the data well. In this adopted geometry, $Ly\alpha$ photons enter the slab normally ($\theta = 0^\circ$), and the H_2 photons that we observe emerge at an angle $\phi = 76^\circ - 79^\circ$ to the normal. This choice is intended to present one approximation of a plausible disk geometry for which we can present detailed results and is not meant to exclude other geometries. In § 5, we present general results for many geometries to demonstrate that our primary result, the temperature of the H_2 gas, is relatively insensitive to geometries, including those that are significantly different from the adopted geometry.

We calculate the relative flux of each transition within a single progression, given an H_2 excitation temperature and column density of the slab, and a separate flux normalization factor for each progression. Models are calculated for temperatures in the range $T = 1000 - 5500$ K, sampled at 100 K intervals, and column densities (in cgs units) of $\log N(H_2) = 15 - 23$, sampled at 0.1 dex intervals. $N(H_2)$ and T are then determined by fitting simultaneously the relative line fluxes for 19 observed progressions. Five lines are not used because they are blended, and one line is not used because it is anomalously weak because of C II wind absorption (see Paper I). The *FUSE* lines, observed six weeks after the STIS observations, are not used in building these models owing to the potential problems with flux variability and cross-calibration between STIS and *FUSE*. The fits have 21 variables: 19 normalization factors, $N(H_2)$, and T . In typical models, 46 of the 140 H_2 line fluxes are sensitive to H_2 line opacity, and hence $N(H_2)$ and T . The other 94 lines, which have only a weak dependence on $N(H_2)$ and T , determine the 19 normalization factors. We calculate $\chi^2_{H_2}$ for only the 46 lines that




FIG. 9.—1–5 σ confidence contours calculated from $\Delta\chi^2_{\nu,H_2}$ for fits to the individual H_2 line fluxes for the thick disk model (see § 3). The star indicates the position of the model with $T = 2500$ K and $\log N(H_2) = 18.5$ (see § 6).

depend on $N(H_2)$ and T . Confidence contours of 1–5 σ , calculated from values of $\Delta\chi^2_{\nu,H_2}$ for 44 degrees of freedom, are shown in Figure 9, with a minimum $\chi^2_{\nu,H_2} = 2.22$ at $\log N(H_2) = 17.9$ and $T = 2600$ K.

Based on this analysis, and additional constraints provided by the $Ly\alpha$ profile analysis described in § 4, we ultimately decide a thick disk model with $T = 2500$ K and $\log N(H_2) = 18.5$ is our best model, which we adopt for the analysis (see § 6.1). Figure 10 compares the observed spectrum compared with a synthetic spectrum computed from this model. The adopted model correctly predicts the fluxes of all strong H_2 lines. Only one observed line flux differs significantly from the model flux. The model predicts a flux of 46×10^{-15} ergs $cm^{-2} s^{-1}$ in the 0–4 $R(1)$ transition at 1333.797 Å. The measured flux in this line, $(7.9 \pm 0.7) \times 10^{-15}$ ergs $cm^{-2} s^{-1}$, is significantly reduced by C II wind absorption, as discussed in Paper I. The 0–3 $P(3)$ transition at 1283.16 Å, with a measured flux of 28.0×10^{-15} ergs $cm^{-2} s^{-1}$, is moderately weaker than the model flux of 34.8×10^{-15} ergs $cm^{-2} s^{-1}$, possibly because of H_2 absorption in the 2–2 $R(15)$ transition, centered 8 km s^{-1} from the 0–3 $P(3)$ transition. Table 3 lists predicted fluxes for Lyman-band H_2 transitions in the *FUSE* bandpass. Future measurements of flux in these transitions would provide a good test for our model. Lines at $\lambda < 1200$ provide particularly good constraints since they are transitions to very low energy levels, and as a result have large opacities.

Neither this model nor the alternative models presented in § 5 rigorously simulate a disk geometry. Because the structure of disks is uncertain near the truncation radii, we do not have an accurate picture of the geometry of the H_2 emission region. Without this a priori knowledge, the thick disk geometry described here is not clearly better or worse than any other oversimplified geometry that we might assume. In § 5, we will show that the temperature of the molecular region is relatively insensitive to geometry, and we will rule out certain geometries that cannot sufficiently explain the observed H_2 spectrum. Since our models make as few assumptions as possible, the primary result of this paper, the temperature of the H_2 gas, may apply to a wide range of other morphologies.

4. RECONSTRUCTED $Ly\alpha$ PROFILE

In this section, we reconstruct the $Ly\alpha$ profile seen by the warm H_2 gas at the 18 pumping wavelengths between 1212




FIG. 10.—Selected regions of the *HST* STIS spectrum of TW Hya. The narrow features are all H_2 lines. The dashed lines show our model fit to the H_2 emission. The line at 1333.8 Å is suppressed by C II in the wind of TW Hya, which places the H_2 interior to the wind.

and 1220 Å. Fluorescent H_2 emission occurs because certain Lyman-band transitions, with lower levels 1–2 eV above the ground state, have wavelength coincidences with the broad $Ly\alpha$ profile. The emission in each progression depends on the $Ly\alpha$ flux at the pumping wavelength of the fluorescent transition, the upward transition probability, the opacity in the pumping transition, and the filling factor of H_2 as seen from the source of the $Ly\alpha$ emission. We reconstruct the incident $Ly\alpha$ profile following a procedure similar to that used by Wood et al. (2002) to analyze fluorescent H_2 emission in the *HST* STIS spectrum of Mira B. A comparison of the reconstructed and observed $Ly\alpha$ profiles provide another constraint on the values of T and $N(H_2)$ that is independent of the constraints provided by the analysis presented in § 3. The analysis in § 3 relied upon using the H_2 flux ratios within each fluorescence progression to constrain T and $N(H_2)$, whereas the $Ly\alpha$ profile reconstruction analysis presented in this section relies only on the total flux from each upper level.

We calculate the optical depth in each H_2 pumping transition by assuming that the absorption line profiles in the model slab are Voigt functions with a Doppler b parameter, which depends on an assumed temperature, and a damping parameter a from Abgrall et al. (1993). The total $Ly\alpha$ flux, F_{abs} , absorbed by a given lower level is then

$$F_{abs} = \eta F_{inc} \int_{\lambda} 1 - e^{-\tau_{\lambda}[T, N(H_2)]} d\lambda, \quad (3)$$

TABLE 3
STRONGEST PREDICTED LYMAN H_2 LINES IN THE *FUSE* BANDPASS

ID	λ_{calc}	F_{mod}	F_{obs}
4–0 $R(3)$	1053.973	1.1	<3
4–0 $P(5)$	1065.593	1.6	<2
4–1 $R(3)$	1101.889	2.4	<10
1–0 $P(5)$	1109.311	1.0	<6
3–1 $P(1)$	1113.877	1.4	<6
4–1 $P(5)$	1113.949	1.4	<6
1–1 $R(3)$	1148.701	5.0	5.5(1.1)
0–1 $R(0)$	1161.693	1.9	13.1(b)
1–1 $P(5)$	1161.814	6.2	13.1(b)
1–1 $R(6)$	1161.949	1.8	13.1(b)
0–1 $R(1)$	1162.170	2.4	13.1(b)
4–1 $R(12)$	1164.596	1.1	<10
2–0 $P(13)$	1165.834	1.0	<6
0–1 $P(2)$	1166.255	2.6	<6
0–1 $P(3)$	1169.751	2.5	<10
4–0 $R(17)$	1176.325	1.0	<15
3–1 $R(12)$	1179.472	1.3	<10
1–1 $P(8)$	1183.309	2.7	<12
2–1 $R(11)$	1185.224	2.8	<12

NOTE.— F_{obs} refers to the observed fluxes and F_{mod} to the model fluxes in units of 10^{-15} ergs cm^{-2} s^{-1} , 1 σ error in parentheses; “(b)” indicates the line is blended with other H_2 lines.

where F_{inc} is the incident $Ly\alpha$ flux per unit wavelength and $\int_{\lambda} 1 - \exp^{-\tau_{\lambda}[T, N(H_2)]} d\lambda$ is the equivalent width of the absorption profile within the slab (see EQW_{slab} in Table 2). Not every absorbed $Ly\alpha$ photon produces an observed H_2 photon. For the most highly excited states, the dissociation probability (P_{dis} in Table 2) can be as large as 42%. Some fraction (R_{unseen} in Table 2) of the emitted photons will be in spectral lines that we did not observe because of line blends, noisy data, or insufficient wavelength coverage. In our plane-parallel slab model, an emerging photon traveling normal to the slab sees a smaller optical depth than a photon traveling nearly parallel to the slab. As a result, the emitted flux in these models is largest for angles normal to the slab.

For comparison with the observed $Ly\alpha$ profile, we compute reconstructed $Ly\alpha$ profiles for the same grid of T and $N(H_2)$ -values used in § 3. Model fluxes at eight pumping transitions, all on the red wing of the $Ly\alpha$ line, are fitted to the observed profile. The observed $Ly\alpha$ flux at other pumping wavelengths is corrupted by interstellar or wind absorption. Two progressions, 3–3 $R(2)$ at 1217.031 Å and 3–3 $P(1)$ at 1217.038 Å, are not included in the fit because their absorption profiles overlap, complicating the analysis. Photoexcitation of H_2 by C III emission via the transition 3–2 $P(4)$ at 1174.923 Å may also add flux to the progression pumped by 3–3 $R(2)$. We judge the suitability of a model using a modified χ^2 analysis to measure the scatter in the red wing of the reconstructed profile. Errors in the $Ly\alpha$ line wing tend to be multiplicative, rather than additive, so we define $\chi^2_{Ly\alpha}$,

$$\chi^2_{Ly\alpha} = \sum \left(\frac{F_{obs}}{\sigma_{obs}} \log \frac{F_{obs}}{F_{mod}} \right)^2, \quad (4)$$

based on standard error propagation techniques from Bevington (1969). Using this formula, an overestimate and an underestimate of the reconstructed $Ly\alpha$ profile by a factor of 10 will result in the same contribution to $\chi^2_{Ly\alpha}$.

Figure 11 shows confidence contours of 1–5 σ calculated from $\Delta\chi^2_{Ly\alpha}$ for fits to the observed $Ly\alpha$ profile, with 5




FIG. 11.—1–5 σ confidence contours calculate from $\Delta\chi^2_{\nu, \text{Ly}\alpha}$ (shaded regions) for our fit of the reconstructed Ly α profile to the red wing of the observed Ly α profile, compared with 1–5 σ confidence contours (solid lines) for fits to the observed H $_2$ fluxes.

degrees of freedom. In the thick disk geometry, a minimum $\chi^2_{\nu, \text{Ly}\alpha} = 2.31$ is obtained when comparing the reconstructed and observed Ly α profiles for $\log N(\text{H}_2) = 18.5$, $T = 2500$ K, and $\eta = 0.25$. The top panel of Figure 12 shows that these parameters yield an excellent fit to the observed profile. These parameters also reproduce the individual line fluxes well (see § 3 and Fig. 9). Since the confidence contours calculated from the Ly α profile analysis place a more stringent constraint on the temperature and column density than do the contours from fitting individual line fluxes (see Fig. 11), we choose on the values of T and $N(\text{H}_2)$ quoted above as our best estimates of these parameters. The model-dependent quantities listed in Table 2 are all calculated using these parameters.

More than 500 other possible upward transitions in the Lyman band have wavelengths between 1210 and 1220 Å. Fluorescent emission in lines from these weaker progressions is below our detection limit. Table 4 identifies undetected progressions with the largest predicted emission line fluxes. The table includes our observational flux limit F_{max} on the strongest line in the progression at wavelength λ_{max} . For any H $_2$ line that is not detected and does not appear in Table 4, we place an upper flux limit of 5×10^{-15} ergs cm $^{-2}$ s $^{-1}$. We then use these upper limits to calculate an upper limit of Ly α at the pumping wavelength to test the self-consistency of our model.

Figure 12 shows three examples of reconstructed Ly α profiles (circles), with upper limits to the Ly α flux at certain wavelengths calculated from upper limits of lines in the undetected progressions (arrows), as described above. The top panel shows the Ly α profile constructed using the parameters for our adopted model [$\log N(\text{H}_2) = 18.5$, $T = 2500$ K, and $\eta = 0.25$]. This profile is smooth, and its wings correspond well with the wings of the observed Ly α profile. The upper limits on the flux of Ly α calculated from undetected progressions are all consistent with the Ly α profile reconstructed from observed progressions. The middle panel shows a Ly α profile calculated from a model with $\log N(\text{H}_2) = 19.0$, $T = 2000$ K, and $\eta = 0.4$, which is inconsistent with the data because of a large amount of scatter in the red wing of the reconstructed Ly α profile. The bottom plot shows a Ly α profile calculated from a model with $\log N(\text{H}_2) = 19.0$, $T = 3000$ K, and $\eta = 0.2$, which is also inconsistent with the data because at 1214.421 Å the predicted Ly α emission required to pump the $v' = 7$, $J' = 4$ level is significantly below the ob-




FIG. 12.—Comparison of the observed (solid line) and calculated (circles) Ly α emission profiles, for models with $\log N(\text{H}_2) = 18.5$, $T = 2500$ K, and $\eta = 0.25$ (top); $\log N(\text{H}_2) = 19.0$, $T = 2000$ K, and $\eta = 0.41$ (middle); and $\log N(\text{H}_2) = 19.0$, $T = 3000$ K, and $\eta = 0.20$ (bottom). The arrows show upper limits to the calculated Ly α emission based on undetected H $_2$ progressions. The top plot is the best fit because of the least scatter in the red wing of Ly α . The bottom model is unacceptable because several undetected progressions ($v' = 7$, $J' = 4$ pumped at 1214.4 Å and $v' = 5$, $J' = 20$ pumped at 1217.7 Å) imply upper limits of Ly α flux that are significantly lower than the flux reconstructed from nearby detected progressions.

servationally constrained flux at 1214.465 Å. Figure 13 is a reproduction of Figure 11, with dotted and diagonal lines added to show the regions of parameter space that are excluded because upper limits on H $_2$ emission would imply a Ly α pumping flux significantly lower than the nearby Ly α pumping flux for a progression of observed H $_2$ lines. Parameter space constraints are shown separately for upper limits on progressions pumped via 7–4 P(5) at 1214.421 Å alone (dots) and via a set of others upward transitions (diagonal lines). The parameter space with $\eta > 1$ (Fig. 13, dashed lines) are unphysical models and are also ruled out.

The three reconstructed Ly α profiles shown in Figure 12 are roughly similar to each other and to most other Ly α profiles that we have reconstructed for other values of T and $N(\text{H}_2)$. Each profile shows a clear depression near Ly α line center that is not nearly as wide as the observed H I absorption. Figure 14 demonstrates that the absorption feature in the reconstructed Ly α profile can be characterized by a column density $\log N(\text{H I}) \approx 18.7$ centered 90 km s $^{-1}$ shortward of line center. This H I between the source of Ly α emission and the warm H $_2$ could be in the base of a wind or at the disk surface. Alternatively, the shape of the Ly α profile could be an intrinsic self-reversal, in which case the intervening H I need not exist. The blue side of the reconstructed Ly α profile is

TABLE 4
UNDETECTED PROGRESSIONS

Pump	λ_{calc} (Å)	A_{ul} (10^{-7} s^{-1})	E'' (eV)	P_{Dis}	F_{max} ($\text{ergs cm}^{-2} \text{ s}^{-1}$)	λ_{max} (Å)	λ_{ref} (Å)	ID _{ref}
1–1 $R(12)$	1212.543	4.6	1.49	5(–5)	3×10^{-15}	1578	1212.425	1–1 $P(11)$
7–4 $P(5)$	1214.421	2.8	2.07	0.19	5×10^{-15}	1582	1214.465	3–1 $R(15)$
5–3 $P(8)$	1218.575	6.6	1.89	0.04	5×10^{-15}	1608	1218.521	2–1 $R(14)$
5–0 $P(20)$	1217.716	5.4	2.39	0.48	2×10^{-15}	1223	1217.904	2–1 $P(13)$
2–2 $P(8)$	1219.154	10.8	1.46	7(–5)	5×10^{-15}	1579	1219.089	0–2 $R(2)$
2–2 $R(9)$	1219.101	12.9	1.56	2(–4)	5×10^{-15}	1572	1219.089	0–2 $R(2)$
5–3 $R(9)$	1219.106	5.4	1.99	0.085	5×10^{-15}	1601	1219.089	0–2 $R(2)$
3–3 $R(1)$	1215.541	0.46	1.48	2(–4)	5×10^{-15}	1596	1215.726	1–2 $R(6)$

enhanced relative to the observed Ly α profile, because wind absorption in our line of sight to the star attenuates the intrinsic emission. This result supports our conclusion from Paper I that, in our line of sight, the H₂ must be photoexcited interior to most of the wind that we detect.

5. ALTERNATIVE GEOMETRIES

In §§ 3 and 4, we described in detail results for a dust-free thick disk geometry, which may not represent the true morphology of warm gas around the star. In this section, we explore the same parameter space for a thin disk geometry, shown schematically in Figure 8, and a cloud geometry, which approximates H₂ emission from a cloud in our line of sight to the star.

The computed H₂ fluxes depend on the angle at which Ly α photons enter the disk, the exit angle of H₂ photons, and the extinction either within the disk or in the ISM. Figures 15 and 16 and Table 5 present the best-fit parameters from fitting the H₂ line opacities for each of three geometries as a function of the exit angle of H₂ and extinction within the disk. In general, while changes in the geometry of the system can mildly change some results, the calculated temperature of the warm gas depends primarily upon the molecular physics and not on the geometry. The calculated column density of H₂ and its physical interpretation, however, do depend on the assumed

geometry. In what follows, the column density represents the amount of H₂ in a line of sight normal to the disk surface, regardless of the path length the Ly α photons traverse through the disk.

5.1. Thin Disk Geometry

A thin disk (see Fig. 8) is characterized here as a disk for which the filling factor of H₂ around the Ly α source is dominated by the surface layers of the disk, rather than by the inner edge of the disk. We approximate this geometry with a model in which the incident Ly α photons enter a plane-parallel slab at a large angle with respect to the disk normal and are reprocessed into H₂ photons that emerge normal to the slab. Because Ly α photons enter the slab at a large angle, the radiation is reprocessed into H₂ emission at a shallow depth, and the emerging H₂ photons see a smaller optical depth in this geometry than in the thick disk geometry. For the thin disk geometry, with incident photons entering the disk at $\theta = 80^\circ$, the best-fit parameters are $T = 2500$ K, $N(\text{H}_2) = 20.0$, but with an unacceptable $\chi^2_{\nu, \text{H}_2} = 7.9$, for fits to H₂ line fluxes. For fits of the reconstructed to the observed Ly α line, the confidence contours move to smaller column densities by a factor of $\sin \theta$. As the angle of incidence asymptotically approaches 90° , the Ly α photons are converted into H₂ photons close to the disk surface. Since the optical depth traversed by the emitted H₂ photons is negligible, the relative line fluxes with




FIG. 13.—Similar contours as in Fig. 11. The dots indicate the parameter space that we rule out because those parameters predict that lines pumped by 7–4 $P(5)$ should be observed, and the diagonal lines indicate the parameter space ruled out because many additional other progressions should be observed but are not detected. Filling factor contours (*dashed lines*) of 0.125–4.0 are also shown. A filling factor above 1 is unphysical. The three models with reconstructed Ly α profiles in Fig. 13 are shown as asterisks.




FIG. 14.—Reconstructed Ly α profile at the wavelengths of the pumping transitions (*diamonds*) for $T = 2500$ K, $\log N(\text{H}_2) = 18.5$, and $\eta = 0.25$ corresponds well to a plausible Ly α profile (*dashed line*), constructed using a single Gaussian profile (*dotted line*) and an absorption feature of $\log N(\text{H I}) = 18.7$ blueshifted 90 km s^{-1} from line center.




FIG. 15.—Best-fit models to H_2 fluxes as a function of exit angle ϕ of H_2 photons, for the thick disk geometry (solid line), thin disk geometry (dashed line), and cloud geometry (dotted line).

each progression differ from predicted branching ratios only because of extinction, which we find to be unlikely (see §§ 2.2 and 5.3).

5.2. Cloud Geometry

We next consider a geometry in which the H_2 is located between us and the source of $Ly\alpha$ photons. In this geometry, the escaping photons must travel through a greater optical depth of H_2 than they would in the other geometries that we considered. As a result, the acceptable contours for fits to H_2 line fluxes move toward smaller column densities relative to the thick or thin disk geometries. Even in this very different geometry, the confidence contours for the $Ly\alpha$ reconstruction fitting do not move much in parameter space, and they restrict the gas temperature to 2000–3000 K. The best-fit parameters in fitting H_2 fluxes for this geometry are $T = 2600$ K, $N(H_2) = 17.8$, with $\chi^2_{\nu, H_2} = 2.8$. The transmission geometry is probably not realistic for TW Hya, but it can be used to set limits on the column of warm H_2 between us and the source of $Ly\alpha$ emission. The absence of any observed H_2 absorption in the $Ly\alpha$ profile (Fig. 3) allows us to place a limit on the column density in this level of $\log N(v'' = 2, J'' = 1) < 14.4$ and a total column density of warm H_2 of $\log N(H_2) < 16.1$. Thus, the detected H_2 fluorescence does not occur in our line of sight to TW Hya.

5.3. Dust Extinction in the Disk and the ISM

In § 2, we presented evidence that the dust extinction along our line of sight to the $Ly\alpha$ emission source is negligible.




FIG. 16.—Best-fit models to H_2 fluxes as a function of extinction within the H_2 slab, for the thick disk geometry (solid line), thin disk geometry (dashed line), and cloud geometry (dotted line). The extinction is for a progression in which the average $Ly\alpha$ photon is absorbed at the midplane of the slab.

Extinction of H_2 emission could occur within the disk if dust is mixed with the warm gas, although Calvet et al. (2002) found that small dust grains are underabundant within 5 AU of the star. We assume an interstellar extinction law from Cardelli et al. (1989), for grains typical of the interstellar medium ($R_V = 3.09$).

For each H_2 progression, the average depth d_{abs} of the initial $Ly\alpha$ absorption depends on the opacity of the lower level of H_2 (see Fig. 17 and Table 2). If a $Ly\alpha$ photon is, on average, absorbed close to the disk surface, then the H_2 lines excited by that photon will not be significantly attenuated by any disk extinction. Thus, the extinction varies for the different progressions. In Figure 16, A_V represents the extinction through half of the slab (not the extinction through the entire slab or the extinction from the star to the observer), which is equal to the largest possible average extinction of H_2 emission when $Ly\alpha$ photons are absorbed uniformly throughout the slab. Disk extinction attenuates both the incident $Ly\alpha$ emission and the outgoing H_2 emission, but not the observed $Ly\alpha$ emission, and thus increases the already large filling factor. In the context of our model, significant disk extinction can be ruled out because the computed filling factors would be much larger than unity.

Interstellar extinction affects the fluxes smoothly across the wavelength region and does not significantly impact the derived filling factor because it attenuates both the observed $Ly\alpha$ emission and the H_2 emission. If we consider only extinction and ignore opacity in the H_2 lines, then from fitting H_2 line fluxes we obtain a minimum $\chi^2_{\nu, H_2} = 6.06$ at $A_V = 1.3$, which is significantly larger than the χ^2_{ν, H_2} for the best-fitting models.

TABLE 5
BEST-FIT PARAMETERS FOR ALTERNATE GEOMETRIES

NUMBER	GEOMETRY ^a				FIT TO H ₂ LINE OPACITIES				FIT TO Ly α PROFILE			
	Type	θ	ϕ	$A_V(D)$	$N(\text{H}_2)$	T	η^b	χ^2_{ν, H_2}	$N(\text{H}_2)$	T	η_{H_2}	$\chi^2_{\nu, \text{Ly}\alpha}$
1.....	Thick	0	89	0	17.0	2200	4.5	1.58	18.7	2400	0.29	0.9
2.....	Thick	0	77	0	17.9	2600	0.41	2.22	18.5	2500	0.25	2.3
3.....	Thick	0	77	0.5	17.5	2700	2.5	2.22	18.8	2800	0.23	9.5
4.....	Thick	0	77	1.0	17.6	2300	16	3.16	18.7	5300	0.09	18
5.....	Thick	0	77	1.5	23.0	1700	0.05	4.21	19.0	5300	0.09	17
6.....	Thick	0	43	0	19.1	2600	0.12	4.15	18.2	2700	0.23	3.0
7.....	Thick	0	0	0	19.5	2500	0.09	4.98	18.4	2600	0.21	3.3
8.....	Thin	80	89	0	16.6	2600	1.10	1.58	18.3	2200	0.31	1.1
9.....	Thin	80	77	0	18.7	2500	0.15	3.97	17.9	2400	0.27	0.8
10.....	Thin	80	43	0	20.2	2500	0.05	6.94	18.0	2400	0.23	0.8
11.....	Thin	80	0	0	20.0	2500	0.05	7.87	18.0	2400	0.22	0.8
12.....	Thin	80	0	0.5	19.2	2500	0.41	7.73	15.3	3300	10 ⁵	29
13.....	Thin	80	0	1.0	17.9	1000	10 ¹³	6.16	19.3	4700	0.03	33
14.....	Thin	80	0	1.5	15.9	4600	10 ¹²	5.86	19.4	4900	0.03	34
15.....	Cloud	0	89	0	16.2	2800	8.2	1.96	18.5	2400	0.38	1.0
16.....	Cloud	0	77	0	17.3	2500	1.2	2.71	18.7	2400	0.26	0.5
17.....	Cloud	0	43	0	17.8	2500	0.52	2.92	18.7	2400	0.25	1.1
18.....	Cloud	0	0	0	17.8	2600	0.64	2.80	18.8	2300	0.33	0.9
19.....	Cloud	0	0	0.5	18.2	2000	12	2.74	18.8	2400	3.32	1.4
20.....	Cloud	0	0	1.0	18.1	1800	400	3.88	18.8	2500	40.4	1.9
21.....	Cloud	0	0	1.5	18.1	1400	10 ⁴	6.07	18.8	2700	400	2.8

^a θ and ϕ are defined in § 3 and Fig. 8.

^b η_{H_2} calculated from Ly α reconstruction for parameters of best fit to H₂ fluxes.

5.4. Adoption of the Thick Disk Geometry

Based on these results, and the results presented in Figures 15–16 and Table 5, we adopt the thick disk geometry (geometry 2 in Table 5) as the preferred geometry to describe the H₂ emission. The thick disk geometry is physically plausible and produces reasonably good fits with self-consistent results, although certain other geometries can produce lower values of χ^2 in fits to H₂ fluxes or in the Ly α fits. Some of those geometries (e.g., geometries 7 and 8 from Table 5)

represent morphologies that are difficult to reconcile with our previous conclusion that the H₂ emission occurs in a disk. In other geometries (e.g., geometries 1 and 11 from Table 5), contours from the H₂ flux fitting do not overlap with the contours from the Ly α fitting as well as they do for the thick disk geometry, i.e., the joint probability is lower (see Fig. 11).

We caution the reader that the thick disk geometry is best only in the context of our restricted set of models. Warm H₂ in the TW Hya star-disk system could in principle have a morphology not considered here.

6. DISCUSSION

6.1. Synthesizing the H₂ and Ly α Modeling Results

Figures 11 and 13, and Table 5, compare results from our analysis of individual H₂ line fluxes with our results from the Ly α profile reconstruction. Based on the overlap region of acceptable models for both the individual H₂ line fluxes and the reconstructed Ly α profile, we conclude that a column density of $\log N(\text{H}_2) = 18.5^{+1.2}_{-0.8}$ is heated to a temperature of 2500^{+700}_{-500} K, with a filling factor of $\eta = 0.25 \pm 0.08$ around TW Hya (2 σ error bars). Table 2 lists the percentage P_{dis} of H₂ that dissociates from the upper level, the H₂ dissociation rate in terms of mass (M_{dis}), and the total flux for each progression (F_{mod}) for our best-fit model of $\log N(\text{H}_2) = 18.5$ and $T = 2500$ K. The notion of a single temperature is a simplistic assumption, but it is sufficient to reproduce our data. The temperature derived here is warmer than the ~ 1100 – 1300 K temperatures seen in the fundamental CO emission by Najita et al. (2003).

The mass column corresponding to $\log N(\text{H}_2) = 18.5$ is 10^{-5} g cm⁻², which is about 10^{-7} times smaller than the mass column predicted to be within 1 AU by D'Alessio et al. (1999). This suggests that H₂ fluorescence probably occurs in a very shallow surface layer of the disk, which could be a




FIG. 17.—Percent of the slab depth that a Ly α photon excites H₂ to two different upper levels (absorption depth d_{abs} in Table 2), at two temperatures. As the column density and temperature of the slab increase, the optical depth in the lower levels increase, and a typical Ly α photon is reprocessed into H₂ at a shallower depth. This transition occurs at different column densities and temperatures for each progression, so that the reconstruction of the Ly α profile is sensitive to these parameters.

warm, extended disk atmosphere. The Ly α fluorescence process cannot be used to detect cold H₂, which most likely represents the bulk of the mass in the disk.

In our models, we assume that H₂ gas is thermalized, so that relative level populations depend only on temperature. This is reasonable because the gas density at the disk surface should be between $n_{\text{H}} = 10^6 - 10^8 \text{ cm}^{-3}$ (Gomez & D'Alessio 2000), which is far larger than the critical density of $n_{\text{H}} = 10^4 \text{ cm}^{-3}$ at which collisions typically dominate over quadrupole radiative de-excitation. Below the critical density, collisions cannot repopulate excited H₂ levels as fast as they radiatively deexcite. Fluorescent progressions pumped out of higher energy states would be weaker than expected from our thermal models, generating scatter that we do not see in our reconstructed Ly α profile. This conclusion confirms the assumption of thermally excited H₂ gas used by Black & van Dishoeck (1987) and Burton, Hollenbach, & Tielens (1990) to model H₂ fluorescence.

6.1.1. Filling Factor of H₂ around the Ly α Source

Figure 13 shows that the filling factor of the H₂ around the Ly α emitting region is $\eta = 0.25 \pm 0.08$. This factor represents the solid angle of the sky subtended by H₂ as seen from the source of Ly α photons near TW Hya, although other geometrical effects may contribute. Neutral hydrogen mixed with the warm H₂ could scatter Ly α photons, increasing the effective path length of the Ly α photons and thereby reducing the calculated filling factor. Any directional dependence of the Ly α emission, and the percentage of Ly α emission visible to us versus the percentage visible to the warm H₂, will also affect the calculated filling factor. A total of 1%–2% of the intrinsic Ly α emission is reprocessed into H₂ emission (see Table 6 and § 4), which places a lower limit of 0.01 on the filling factor.

For any flared disk geometry, the filling factor can be converted into a disk height by comparing the surface areas of a cylinder and a sphere. For $\eta = 0.25$, the implied geometrical disk height from the midplane to the surface at distance R from the center of the star is $H = 0.25(R/1 \text{ AU}) \text{ AU}$, compared with predicted disk heights from D'Alessio (2001) of about 0.15 AU at a distance of 1 AU from the star.

6.1.2. Relationship between UV and IR H₂ Emission

Emission in the 1–0 $S(1)$ rovibrational transition at 2.1218 μm has been detected from four T Tauri stars (Bary et al. 2003). The narrow width and absence of any velocity shift with respect to the stars suggests a disk origin of this emission. The flux in this line from an optically thin emission region

$$F = \frac{N(1,3)\pi r^2 A_{ul} hc}{4\pi d^2 \lambda}, \quad (5)$$

where $N(1, 3)$ is the H₂ column density of the disk in the $v'' = 1$, $J'' = 3$ level, πr^2 is the projected surface area of the emission region, and $A_{ul} = 3.47 \times 10^{-7} \text{ s}^{-1}$. If the 1–0 $S(1)$ line flux of $1 \times 10^{-15} \text{ ergs cm}^{-2} \text{ s}^{-1}$ measured by Weintraub, Kastner, & Bary (2000) comes from a heated disk surface layer with $N(\text{H}_2) = 18.5$ and $T = 2500 \text{ K}$, then the 1–0 $S(1)$ emission must extend out to at least 5 AU for a face-on disk. This corresponds to an angular extent of $\pm 0.7''$, which could have been resolved in our *HST* observations but was not. Thus, it seems likely that at least some of the 1–0 $S(1)$ emission comes from H₂ gas that is not fluoresced by Ly α . Some of the emission in the 1–0 $S(1)$ line may be produced

deeper in the disk or farther from the star, as a result of X-ray ionization and subsequent collisional excitation of H₂ by nonthermal electrons (Weintraub et al. 2000).

6.1.3. H₂ Dissociation Rate

From the flux in observed transitions, we calculate a dissociation rate of $9.8 \times 10^{-11} M_{\odot} \text{ yr}^{-1}$ for H₂ in the disk. Forward modeling of undetected Lyman and Werner-band transitions excited by Ly α photons increases the calculated dissociation rate to $1.6 \times 10^{-10} M_{\odot} \text{ yr}^{-1}$. Depending on the timescale for collisional de-excitation of H₂, the actual dissociation rate could be much higher as a result of multiple pumping (pumping from lower rovibrational levels of H₂ that are populated by the initial pump and subsequent fluorescence—see Shull 1978 for details), which can populate highly excited rovibrational levels in the B electronic state that have dissociation probabilities as high as 50%.

The H₂ dissociation rate we estimate is within a factor of 10 of TW Hya's mass accretion rate (see § 2.1), meaning that the dissociation provided by Ly α fluorescence may be an important process in the accretion of hydrogen onto the star. The neutral H released by dissociation will cause the incident Ly α to scatter, increasing the probability of absorption in the pumping transitions and therefore decreasing the required filling factor. The neutral H could also be sufficiently optically thick to produce the self-reversal of the Ly α profile that irradiates the disk (see Fig. 14).

6.2. FUV Radiation Field

The FUV radiation field incident on the accretion disk can significantly affect the chemistry and temperature structure of the disk. In Table 6 we list the flux for spectral features in the FUV spectrum of TW Hya. The observed Ly α line contributes 67% of the observed FUV flux of $6.5 \times 10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1}$ between 1170 and 1700 Å. From the reconstructed Ly α profile, we infer an intrinsic flux of $(8-16) \times 10^{-11} \text{ ergs cm}^{-2} \text{ s}^{-1}$, depending on the correction for the central absorption. With this correction, Ly α emission contributes 80%–90% of the FUV flux of TW Hya, of which 1%–2% is reprocessed into H₂ emission. The continuum accounts for about $6 \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$, or about 4% of the total emission, including the estimated intrinsic Ly α flux. The remaining flux occurs in other emission lines such as the C IV 1550 Å doublet. The UV flux below 1170 Å measured by *FUSE* adds at most 2% additional flux to the strength of the radiation field. The UV radiation between 1700 and 2000 Å may add to the radiation field, primarily because of emission in lines such as Si II λ 1817 and C III λ 1909 but will have little effect on the excitation of molecules and atoms in the disk. The estimated total emission after including Ly α flux that is not observable is approximately $10^{-3} L_{\odot}$, assuming isotropic emission and a distance of 56 pc.

The properties of the disk, both in the PDR-like disk atmosphere and in the cooler outer regions, will be partially controlled by the Ly α emission, as evidenced by the fluorescent H₂ emission. Not only does Ly α dominate the FUV flux of TW Hya, but it can also scatter off neutral H toward the interior of the disk, allowing it to penetrate the disk further than continuum UV radiation. Ly α can dissociate molecules such as H₂ and H₂O and can ionize Si and C. Bergin et al. (2003) explain that the enhancement of CN relative to HCN occurs in disks because Ly α radiation can photodissociate HCN, whereas CN can be dissociated only by radiation with $\lambda < 1150 \text{ Å}$.

TABLE 6
FLUXES OF STRONG LINES

ID (Å)	λ_{lab}	Flux ^a (10^{-12} ergs cm ⁻² s ⁻¹)
Total flux.....	1170–1700	65
Ly α ^b	1215.67	44.7 ^{de}
Ly α ^c	1215.67	80–160
Continuum.....	1170–1700	6.0 ^f
H ₂ (obs).....	<1700	1.94 ^j
H ₂ (mod).....	<1700	2.20 ^k
C IV.....	1548	1.86 ^f
He II.....	1641	1.35
C IV.....	1551	0.98
C III.....	977	0.44 ^{dh}
C III.....	1175	0.42
C III.....	1175	0.35 ⁱ
O I.....	1306	0.37
O VI.....	1032	0.31 ^g
N V.....	1238	0.30 ^f
O I.....	1305	0.30 ^d
O I.....	1302	0.28 ^{de}
O VI.....	1038	0.15 ^g
C II.....	1336	0.23 ^d
N V.....	1243	0.12
C II.....	1335	0.11 ^{de}
Si IV.....	1393	0.076 ^f
Si IV.....	1402	0.034 ^f
O III.....	1666	0.023

^a These fluxes are more accurate than those in Table 3.

^b Observed Ly α flux.

^c Flux estimated from reconstructed Ly α profile.

^d Wind absorption attenuates some blue emission.

^e IS/CS absorption absorbs some emission.

^f Flux from blended H₂ lines subtracted.

^g Average flux from *FUSE* LiF1A, LiF2B channels.

^h Average flux from *FUSE* SiC2A, SiC1B channels.

ⁱ Average flux from *FUSE* LiF2A, LiF1B channels.

^j Observed H₂ flux.

^k Total H₂ flux, including weak fluxes from model.

6.3. Heating Mechanisms

We have found evidence for a warm surface layer of H₂ on the disk around TW Hya. From the lack of spatial extent in the STIS echelle cross-dispersion direction, the heated surface is located within 2 AU of the star. Models of dust in disks of CTTSs typically predict $T \approx 1000$ K only within 0.1 AU of the star (D'Alessio, Calvet, & Hartmann 2001; Chiang & Goldreich 1997). The FUV radiation field at 1 AU from the star is approximately $10^7 G_0$, where the Habing field $G_0 = 1.6 \times 10^{-3}$ ergs cm⁻² s⁻¹ (the local interstellar field is $1.7 G_0$), which is stronger than the FUV radiation field incident upon most PDRs (10^3 – $10^6 G_0$). Hollenbach, Yorke, & Johnstone (2000) predicted from models of photodissociation regions that a warm atmosphere-like structure, heated by UV radiation, should surround the inner regions of a disk, with a temperature between 500 and 3000 K.

Because of the absence of micron-sized dust grains within 5 AU of TW Hya (Calvet et al. 2002), a large percentage of the FUV radiation may be deposited into the gas. In their analysis of the H₂ fluorescence from Mira B, Wood et al. (2002) found that the temperature increases sharply with radius from the UV source and that the Ly α fluorescence process is a significant heating source for the gas. The energy deposited into the warm gas near TW Hya as a result of Ly α photo-excitation and subsequent fluorescence is 1.4×10^{29} ergs s⁻¹,

which corresponds to roughly 0.01 eV s⁻¹ per molecule of H₂. The strong X-ray luminosity of TW Hya, $L_X = 3.6 \times 10^{-4} L_\odot$ between 0.45 and 6.0 keV (Kastner et al. 2002), could heat the disk surfaces by an additional 500 K (Igea & Glassgold 1999; Glassgold & Najita 2001; Fromang, Terquem, & Balbus 2002) by ionizing atoms to produce a large reservoir of energetic nonthermal electrons (e.g., Maloney, Hollenbach, & Tielens 1996; Yan & Dalgarno 1997). X-rays can also penetrate deeper into the gas than UV radiation and, consequently, may heat gas deeper in the disk than the UV radiation.

Certain observed fluorescence progressions are pumped from lower rovibrational levels with large energies, such as $v'' = 3$, $J'' = 25$ ($E'' = 4.2$ eV) and $v'' = 5$, $J'' = 18$ ($E'' = 3.8$ eV). The measured flux in these progressions, combined with the observed flux at the pumping wavelengths, requires that of order 1% of the H₂ resides in these levels. This population is too large to be thermally populated at any reasonable temperature for the H₂ gas. Therefore, nonthermal processes must be populating these levels. Although the fluorescence process can populate high vibrational levels in the ground electronic state, fluorescence cannot directly populate high rotational levels, and no direct paths exist to the levels $v'' = 3$, $J'' = 25$ and $v'' = 5$, $J'' = 18$. H₂ can form in highly excited states (Kokoouline, Greene, & Esry 2001; Strasser et al. 2001) via dissociative recombination of H₃⁺. Given a dense medium and a large ionizing flux, then H₂⁺ may be present in large enough quantities to react with H₂ to produce H₃⁺ in highly excited states. More attempts to detect H₃⁺ would be useful.

7. SUMMARY

We have modeled a plane-parallel slab of warm H₂ (see Fig. 1), which represents the surface of a protoplanetary disk irradiated by UV flux from the star. Our analysis of the ultraviolet spectrum of TW Hya obtained with the STIS instrument on *HST* and with *FUSE* leads to the following conclusions:

1. The FUV continuum rises shortward of 1700 Å, which is indicative of an H₂ dissociation continuum, although it could also be produced by H₂ fluorescence resulting from FUV continuum pumping or an additional accretion or activity component. The FUV continuum is not well explained by simple models of a pure hydrogen slab, which are commonly invoked to analyze the excess NUV continuum.

2. The extinction toward TW Hya is negligible, based on the hydrogen column density in our line of sight and assuming an interstellar gas-to-dust ratio.

3. Self-absorption of Lyman-band transitions involving low excitation energy levels weakens the flux in these lines. We model this effect by simulating Ly α emission entering a plane-parallel atmosphere and pumping the H₂.

4. Using the observed H₂ fluxes and our fluorescence models, we reconstruct the Ly α line profile incident upon the warm molecular layer and compare it with the observed Ly α profile, for a range of assumed temperatures and column densities. Undetected progressions rule out a large region of parameter space in this model. The reconstructed Ly α profile is similar to the observed profile in the wings but shows a much narrower absorption feature than is observed. This narrow absorption component in the reconstructed profile could be a self-reversal or a component of the wind between the source of Ly α emission and the warm molecular region.

5. Our models indicate that a molecular layer with a kinetic temperature of 2500_{-500}^{+700} K and a column density of

$\log N(\text{H}_2) = 18.5^{+1.2}_{-0.8}$ (2σ error bars) absorbs $\text{Ly}\alpha$ radiation in the surface layer and inner edge of the disk within 2 AU of the central star. The $\text{Ly}\alpha$ pumping leads to a small H_2 dissociation rate and does not cause significant disk dissipation. The filling factor of the warm H_2 around TW Hya is $\eta = 0.25 \pm 0.08$, although significant uncertainties in the geometry of the fluorescent H_2 weaken our confidence in the large filling factor.

6. The warm H_2 most likely resides in a warm surface layer of the disk. This surface layer may be analogous to a classic PDR, although in this case the FUV radiation field is dominated by emission lines rather than a continuum. In particular, $\text{Ly}\alpha$ comprises about 85% of the FUV radiation field below 2000 Å, and it controls the excitation and ionization of the disk surface.

7. Some of the observed H_2 lines are pumped from high rotational levels that cannot be excited thermally or by fluo-

rescence. Formation of rotationally excited H_2 by reactions with H_3^+ may explain these emission lines.

This research is supported by NASA grant S-56500-D to the University of Colorado and NIST, and by a grant to the University of Colorado of NASA funds through the Johns Hopkins University. We thank the referee for thorough and valuable comments. The referee especially helped to significantly improve the clarity of the paper. We also thank Chris Greene and Ben McCall for valuable discussions about H_2 formation, and Phil Maloney for insightful comments about PDRs.

REFERENCES

- Abgrall, H., Roueff, E., & Drira, I. 2000, *A&AS*, 141, 297
 Abgrall, H., Roueff, E., Launay, F., Roncin, J. Y., & Subtil, J. L. 1993, *A&AS*, 101, 273
 Agnor, C. B., & Ward, W. R. 2002, *ApJ*, 567, 579
 Aikawa, Y., van Zadelhoff, G. J., van Dishoeck, E. F., & Herbst, E. 2002, *A&A*, 386, 622
 Alencar, S. H. P., & Batalha, C. 2002, *ApJ*, 571, 378
 Ardila, D. R., Basri, G., Walter, F. M., Valenti, J. A., & Johns-Krull, C. M. 2002a, *ApJ*, 566, 1100
 ———. 2002b, *ApJ*, 567, 1013
 Bary, J. S., Weintraub, D. A., & Kastner, J. H. 2003, *ApJ*, 586, 1136
 Bergin, T., Calvet, N., D'Alessio, P., & Herczeg, G. J. 2003, *ApJ*, 591, 159
 Bevington, P. R. 1969, *Data Reduction and Error Analysis for the Physical Sciences* (New York: McGraw-Hill Book)
 Black, J. H., & van Dishoeck, E. F. 1987, *ApJ*, 322, 412
 Bohlin, R. C., Savage, B. D., & Drake, J. F. 1978, *ApJ*, 224, 132
 Burton, M. G., Hollenbach, D. J., & Tielens, A. G. G. M. 1990, *ApJ*, 365, 620
 Calvet, N., D'Alessio, P., Hartmann, L., Wilner, D., Walsh, A., & Sitko, M. 2002, *ApJ*, 568, 1008
 Calvet, N., Patino, A., Magrid, G. C., & D'Alessio, P. 1991, *ApJ*, 380, 617
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Carr, J. S., Tokunaga, A. T., Najita, J., Shi, F. H., & Glassgold, A. E. 1993, *ApJ*, 411, L37
 Casali, M. M., & Eiroa, C. 1996, *A&A*, 306, 427
 Chiang, E. I., & Goldreich, P. 1997, *ApJ*, 490, 368
 Curiel, S., Raymond, J. C., Wolfire, M., Hartigan, O., Morse, J., Schwartz, R. D., & Nissen, P. 1995, *ApJ*, 453, 322
 D'Alessio, P. 2001, in *ASP Conf. Ser. 244, Young Stars Near Earth: Progress and Prospects*, ed. R. Jayawardhana & T. Greene (San Francisco: ASP), 239
 D'Alessio, P., Calvet, N., & Hartmann, L. 2001, *ApJ*, 553, 321
 D'Alessio, P., Calvet, N., Hartmann, L., Lizano, S., & Canto, J. 1999, *ApJ*, 527, 893
 Fromang, S., Terquem, C., & Balbus, S. A. 2002, *MNRAS*, 329, 18
 Gomez, J. F., & D'Alessio, P. 2000, *ApJ*, 535, 943
 Gomez de Castro, A. I., & Lamzin, S. A. 1999, *MNRAS*, 304, L41
 Glassgold, A., & Najita, J. 2001, in *ASP Conf. Ser. 244, Young Stars Near Earth: Progress and Prospects*, ed. R. Jayawardhana & T. Greene (San Francisco: ASP), 251
 Goldreich, P., & Tremaine, S. 1979, *ApJ*, 233, 857
 Gullbring, E., Hartmann, L., Briceno, C., & Calvet, N. 1998, *ApJ*, 492, 323
 Hartigan, P., Kenyon, S. J., Hartmann, L., Strom, S. E., Edwards, S., Welty, A. D., & Stauffer, J. 1991, *ApJ*, 382, 617
 Herczeg, G. J., Linsky, J. L., Valenti, J. A., & Johns-Krull, C. M. 2002, *ApJ*, 572, 310 (Paper I)
 Hollenbach, D. J., Yorke, H. W., & Johnstone, D. 2000, in *Protostars & Planets IV*, ed. V. Mannings, A. P. Boss, & S. S. Russell (Tucson: Univ. Arizona Press), 401
 Igea, J., & Glassgold, A. E. 1999, *ApJ*, 518, 848
 Johns-Krull, C. M., & Valenti, J. A. 2001a, *ApJ*, 561, 1060
 Johns-Krull, C. M., & Gafford, A. D. 2002, *ApJ*, 573, 685
 Kastner, J. H., Huenemoerder, D. P., Schulz, N. S., Canizares, C. R., & Weintraub, D. A. 2002, *ApJ*, 567, 434
 Kastner, J. H., Huenemoerder, D. P., Schulz, N. S., & Weintraub, D. A. 1999, *ApJ*, 525, 837
 Kenyon, S. J., & Hartmann, L. 1995, *ApJS*, 101, 117
 Kokouline, V., Greene, C. H., & Esry, B. D. 2001, *Nature*, 412, 891
 Krist, J. E., Stapelfeldt, K. R., Ménard, F., Padgett, D. L., & Burrows, C. J. 2000, *ApJ*, 538, 793
 Lissauer, J. J. 1993, *ARA&A*, 31, 129
 Liu, W., & Dalgarno, A. 1996, *ApJ*, 467, 446
 Lynden-Bell, D., & Pringle, J. E. 1974, *MNRAS*, 168, 603
 Maloney, P. R., Hollenbach, D. J., & Tielens, A. G. G. M. 1996, *ApJ*, 466, 561
 Mamajek, E. E., Lawson, W. A., & Feigelson, E. D. 2000, *ApJ*, 544, 356
 Muzerolle, J., Calvet, N., Briceno, C., Hartmann, L., & Hillenbrand, L. 2000, *ApJ*, 535, 47
 Najita, J., Carr, J. S., Glassgold, A. E., Shu, F. H., & Tokunaga, A. T. 1996, *ApJ*, 462, 919
 Najita, J., Carr, J. S., & Mathieu, R. D. 2003, *ApJ*, 589, 931
 Rachford, B. L., et al. 2002, *ApJ*, 577, 221
 Raymond, J. C., Blair, W. P., & Long, K. S. 1997, *ApJ*, 489, 314
 Redfield, S., & Linsky, J. L. 2000, *ApJ*, 534, 825
 Richter, M. J., Jaffe, D. T., Blake, G. A., & Lacy, J. H. 2001, *BAAS*, 199, 6005
 Roberge, A., et al. 2001, *ApJ*, 551, L97
 Saucedo, J., Calvet, N., Hartmann, L., & Raymond, J. 2003, *ApJ*, 591, 275
 Savage, B. D., Bohlin, R. C., Drake, J. F., & Budich, W. 1977, *ApJ*, 216, 291
 Schwartz, R. D. 1983, *ApJ*, 268, L37
 Sheret, I., Ramsay Howat, S. K., & Dent, W. R. F. 2003, *MNRAS*, 343, L65
 Shu, F., Najita, J., Ostriker, E., Wilkin, F., Ruden, S., & Lizano, S. 1994, *ApJ*, 429, 781
 Shull, J. M. 1978, *ApJ*, 219, 877
 Strasser, D., et al. 2001, *Phys. Rev. Lett.*, 86, 779
 Thi, W. F., et al. 2001, *ApJ*, 561, 1074
 Trilling, D. E., Koerner, D. W., Barnes, J. W., Ftaclas, C., & Brown, R. H. 2001, *ApJ*, 552, 151
 Valenti, J. A., Basri, G., & Johns, C. M. 1993, *ApJ*, 106, 2024
 Walter, F. M., et al. 2003, *AJ*, 126, 3076
 Webb, R. A., Zuckerman, B., Patience, J., White, R. J., Schwartz, M. J., McCarthy, C., & Platais, I. 1999, *ApJ*, 512, L63
 Weinberger, A. J., et al. 2002, *ApJ*, 566, 409
 Weintraub, D. A., Kastner, J. H., & Bary, J. S. 2000, *ApJ*, 541, 767
 White, R. J., & Ghez, A. M. 2001, *ApJ*, 556, 265
 Wilner, D. J., Ho, P. T. P., Kastner, J. H., & Rodriguez, L. F. 2000, *ApJ*, 534, L101
 Wood, B. E., Karovska, M., & Raymond, J. C. 2002, *ApJ*, 575, 1057
 Yan, M., & Dalgarno, A. 1997, *ApJ*, 481, 296
 Zuckerman, B. 2001, *ARA&A*, 39, 549
 Zuckerman, B., Forveille, T., & Kastner, J. H. 1995, *Nature*, 373, 494